

Physical and shift work in pregnancy

Occupational
aspects of
management

A national guideline



Royal College
of Physicians
Setting higher medical standards

NHS
Plus

Physical and shift work in pregnancy

Occupational aspects of
management

A national guideline

2009



Royal College
of Physicians
Setting higher medical standards

NHS
Plus

Acknowledgements

We are indebted to the research librarians, Lesley Vickers and Susan Paterson, for undertaking the literature searches and for retrieving the papers. We would also like to thank Barbara Smiley for her administrative support towards the end of the project.

NHS Plus

The NHS Plus Project aims to improve the quality and delivery of occupational health services to NHS staff and in turn increase the availability of NHS Plus services to small and medium employers. In addition to commissioning the Occupational Health Clinical Effectiveness Unit to produce evidence-based guidelines and conduct national audits, the Project has work strands to improve the delivery of services, provide an improved trading model and improve the strategic leadership of occupational health services in the NHS.

The Royal College of Physicians

The Royal College of Physicians plays a leading role in the delivery of high-quality patient care by setting standards of medical practice and promoting clinical excellence. We provide physicians in the United Kingdom and overseas with education, training and support throughout their careers. As an independent body representing over 20,000 Fellows and Members worldwide, we advise and work with government, the public, patients and other professions to improve health and healthcare.

Occupational Health Clinical Effectiveness Unit

The Occupational Health Clinical Effectiveness Unit at the Royal College of Physicians aims to measure and raise standards, and to reduce variability, of occupational healthcare through the development of evidence-based guidelines and conduct of national clinical audits.

Faculty of Occupational Medicine

Our aim is for healthy working lives through:

- elimination of preventable injury and illness caused or aggravated by work
- maximising people's opportunities to benefit from healthy and rewarding work while not putting themselves or others at unreasonable risk
- access for everyone to advice from a competent occupational physician as part of comprehensive occupational health and safety services.

Citation for this document

NHS Plus, Royal College of Physicians, Faculty of Occupational Medicine. *Physical and shift work in pregnancy: occupational aspects of management. A national guideline*. London: RCP, 2009.

Copyright

All rights reserved. No part of this publication may be reproduced in any form (including photocopying or storing it in any medium by electronic means and whether or not transiently or incidentally to some other use of this publication) without the written permission of the copyright owner. Applications for the copyright owner's written permission to reproduce any part of this publication should be addressed to the publisher.

Copyright © 2009
Royal College of Physicians

ISBN 978-1-86016-353-1

Review date: 2014

Royal College of Physicians of London
11 St Andrews Place, London NW1 4LE

www.rcplondon.ac.uk

Registered Charity No 210508

Cover design: WLG Design

Typeset by Dan-Set Graphics,
Telford, Shropshire

Printed by The Lavenham Press Ltd,
Sudbury, Suffolk

Contents

	Acknowledgements	ii
	Guideline Development Group	v
	Executive summary	vii
	Glossary	ix
1	Introduction	1
	Aim	1
	Background	1
	Epidemiology of adverse pregnancy outcomes	2
	Possible biological mechanisms	4
	Evidence of variation in practice: relevance to occupational medicine	5
2	Methodology of the evidence review	7
	Key questions	7
	Literature search	8
	Critical appraisal	11
	Assessing the evidence	11
	Development of recommendations	12
	Limitations of the evidence base	12
3	Findings and recommendations	15
	Synthesis of the evidence and recommendations	15
	1 Manual handling, lifting, heavy physical work	15
	(a) Heavy physical effort	15
	Recommendations	16
	(b) Lifting	17
	Recommendations	18
	2 Prolonged standing	18
	Recommendations	19
	3 Working hours	20
	Recommendations	21
	4 Shift work	21
	5 Fatigue score	22
4	Future research	23
	Recommendations for research	23
5	Audit criteria	24
	Evidence tables	
	1 Summary of papers included as evidence	25
	2 Preterm birth/threatened preterm labour	28
	3 Spontaneous miscarriage/perinatal mortality	37

Physical and shift work in pregnancy: occupational aspects of management

4 Low birth weight	43
5 Pre-eclampsia	47
6 Small for gestational age/intra-uterine growth restriction	49
7 Sick leave/sick certification	51
Appendices	
1 Search strategy	54
2 Critical appraisal form	56
3 SIGN grading	58
4 Grading system for recommendations	59
5 Score per paper	60
References	79

Guideline Development Group

A multidisciplinary Guideline Development Group (GDG) was established at the outset of the project. Each member brought their own expertise, and both a personal and a professional perspective. The group included obstetric and midwifery clinicians, occupational health practitioners and human resources professionals.

GDG Leader

Dr Nadia Sheikh, Specialist Registrar in Occupational Medicine,
Royal Free Hampstead NHS Trust, London

Guideline Editor

Dr Julia Smedley, Lead Consultant and Head of Occupational Health,
Southampton University Hospitals NHS Trust

Director of evidence-based guideline project, NHS Plus

Dr Ira Madan, Consultant and Senior Lecturer in Occupational Medicine,
Guys and St Thomas' NHS Foundation Trust and King's College London

External assessors

Professor Keith Palmer, Professor of Occupational Medicine,
Medical Research Council

Professor Khalid S Khan, Professor of Obstetrics-Gynaecology and Clinical Epidemiology,
Birmingham Women's Hospital

Dr Susan Bewley, Consultant Obstetrician,
Guy's and St Thomas' NHS Foundation Trust

Guideline Development Group Members

Dr Paul Grime, Director of Occupational Health, Royal Free Hampstead NHS Trust

Dr Laila Kapadia, Occupational Health Physician, St Georges Healthcare NHS Trust

Ms Amanda Mansfield, Consultant Midwife, Royal Free Hampstead NHS Trust

Ms Sharon O'Byrne, Assistant Director of Human Resources,
Royal Free Hampstead NHS Trust

Ms Chris Perry, Domestic Services Manager, Royal Free Hampstead NHS Trust

Ms Lesley Purdie, Senior Back Care Advisor, HCA International

Miss Elaine Scott, Consultant Obstetrician, Royal Free Hampstead NHS Trust

Ms Maggi Smith, Senior Occupational Health Nurse, Royal Free Hampstead NHS Trust

Conflicts of interest: none declared.

Executive summary

The purpose of these guidelines is to enable consistent evidence-based advice to be given in relation to pregnant women who are exposed to hazards at work. The guidance is intended for use by health professionals who are advising working women and their employers. The document is focused on a number of specific hazards, namely lifting/manual handling and heavy physical work, prolonged standing, long working hours and shift work (including night shifts). The evidence considered applies to women who are well during their pregnancy; those with pregnancy-related complications need to be considered on an individual basis.

The production of this full guideline and the accompanying short guidance documents for employers, employees and health professionals was overseen by a steering committee, the Guideline Development Group (GDG). The GDG provided multidisciplinary, practitioner-based input both to the process and the emerging messages for practice, and aimed to reduce author bias.

The GDG agreed the scope of the guideline in advance, including the key questions to be addressed. Subsequently a literature search, based upon the key questions, was undertaken by two librarians at the Department of Health. This yielded 270 abstracts, from which selected papers were retrieved according to pre-determined inclusion criteria. Thirty-seven relevant papers met the inclusion criteria and were critically appraised by members of the GDG.

Each paper was appraised using the Scottish Intercollegiate Guidelines Network (SIGN) system. Papers were assigned a SIGN grade, which took account of the type of study and the methodological rigour. In order to increase the consistency and transparency of assignment of the SIGN grading, a standardised scoring system was devised and used by appraisers to assess important methodological limitations such as response rates, bias and confounding. Based upon the critical appraisals, the volume and quality of evidence was synthesised using the SIGN method and a series of graded recommendations were made.

The results show that there is consistent evidence of risk for some adverse pregnancy outcomes in relation to lifting and heavy work, prolonged standing and long working hours, but the level of risk is small. There was reasonable evidence that shift work is not associated with a high risk of adverse pregnancy outcome, although there was some conflicting evidence and further research in this area would be helpful. Apart from one high quality systematic review (including a meta-analysis and rated 1++ according to the SIGN convention), quality ratings were confined to SIGN category 2++ or below. This was partly due to a reliance on observational studies, which are common in occupational epidemiology. Whilst the number of research papers was large for some adverse pregnancy outcomes it was small for others, in particular pre-eclampsia. There were conflicting results for some outcome-exposure combinations (particularly preterm birth with heavy physical work, and preterm birth with shift or night work). Because of these uncertainties, and to adopt a precautionary approach, recommendations have been made for employers to reduce exposure to lifting, heavy physical work and prolonged standing for pregnant employees. Due to a lack of detailed evidence, it was not possible to specify either the level of exposure at which adjustments should be applicable or the stage of pregnancy at which they should be implemented for some of the outcome-exposure combinations. None of the risks were sufficiently strong to justify mandatory exclusion of pregnant women from work for any of the exposures that were considered.

Glossary

Birth weight:

Low birth weight Birth weight <2500 g

Very low birth weight Birth weight <1500 g

Extremely low birth weight Birth weight <1000 g

Dubowitz score A method of clinical assessment in the newborn from birth until five days old that includes neurological criteria for the infant's maturity and other physical criteria to determine gestational age.

Gestation The time from conception to birth. The duration of gestation is measured from the first day of the last normal menstrual period.

Hypertension Diastolic BP ≥ 110 mmHg on any one occasion, or diastolic BP ≥ 90 mmHg on two or more consecutive occasions more than four hours apart.

Systolic BP 30 mmHg above earliest recorded in pregnancy or diastolic increase of 15–25 mmHg.

Moderate hypertension (2 readings of >100 mmHg diastolic blood pressure) with significant proteinuria and at least 2 signs or symptoms of imminent eclampsia, will include many women with severe pre-eclamptic toxæmia (PET).

Severe hypertension A diastolic blood pressure >110 mmHg on two occasions, *or* systolic blood pressure >170 mmHg on two occasions.

Intra uterine growth restriction (IUGR) This is a term often used interchangeably with the term SGA (small for gestational age). However, IUGR strictly refers to babies that have failed to reach their genetic growth potential during pregnancy. They are frequently but not always SGA. SGA is variably defined as a baby/fetus with measurements less than the 3rd centile, 5th centile or 10th centile.

Late fetal loss A death occurring between 22 weeks + 0 days and 23 weeks + 6 days. If gestation is not known or not sure, all births of at least 300 g are reported. Late fetal loss and stillbirth are distinguished by gestational age at the time of delivery, which is not necessarily the time of death (CESDI).

Miscarriage The spontaneous loss of a pregnancy before 24 completed weeks of gestation.

Pre-eclampsia Pregnancy-induced hypertension in association with proteinuria (>0.3 g/24 hours) + oedema and virtually any organ system may be affected (eg cardio-vascular, renal, central nervous system, hepatic, coagulation and placenta).

Severe pre-eclampsia Severe hypertension associated with significant proteinuria (at least 1 g/l).

Prelabour rupture of membranes (PROM) Rupture of membranes with leakage of amniotic fluid in the absence of spontaneous uterine activity. Previously often described as premature rupture of membranes.

Preterm prelabour rupture of membranes PROM before 37 completed weeks of gestation.

Preterm labour Onset of labour before 37 completed weeks of gestation.

Preterm delivery Delivery of the fetus before 37 completed weeks of gestation.

Prolonged standing Continuous standing for more than three hours.

Small for gestational age (SGA) See IUGR.

Stillbirth (England and Wales) A child that has issued forth from its mother after the 24th week of pregnancy and which did not at any time, after being completely expelled from its mother, breathe or show any other signs of life.

Stillbirth (Northern Ireland) The complete expulsion from its mother after the 24th week of pregnancy of a child, which did not at any time after being completely expelled or extracted, breathe or show any other evidence of life.

(Note: before 1992, it was 28 weeks' gestation.)

The WHO has chosen 22 weeks rather than 24, but because of the difficulty in date accuracy they use fetal weight: death of a fetus at least 500 g before expulsion/extraction from its mother.

Threatened preterm labour Contractions or cervical change, before 37 weeks' gestation.

Trimester of pregnancy Pregnancy is traditionally divided into three time periods. The first trimester is from conception until 12 completed weeks' gestation. The third trimester is from 28 weeks until delivery.

1 Introduction

Aim

The broad aim of these guidelines is to provide evidence-based advice for health professionals advising pregnant women and employers with respect to identifiable hazards in the workplace. The scope of the document comprises the following specific hazards:

- (1) Manual handling/lifting/heavy physical work
- (2) Prolonged standing
- (3) Long weekly working hours (typically >40 per week)
- (4) Shift work including night work.

Background

Women at work

There has been a world-wide increase in the prevalence of women working during pregnancy.⁵⁹ In 1995 in the UK, 66% of women of working age were in employment. This figure had risen to 69% by 2000²⁶ and continued to rise. In 2004, seasonally adjusted figures for the spring quarter showed that 45% of the employed population were women. The majority of these (77%) were of child-bearing age (18–49 years old).^{99,109} Over half of women who are pregnant in the UK are in paid employment during their pregnancy and 37% were still working within six weeks of delivery.⁶¹ Every year around 350,000 women continue to work during their pregnancy and, of these, 69% return to work soon after giving birth (Tommy's website: www.tommys.org source: Department of Trade and Industry).

The majority of women remain well throughout their pregnancy. Pregnancy should not be regarded either as an illness or, generally, as a contraindication to work. Indeed there is some evidence of a beneficial effect of work on pregnancy. It has been suggested that the 'reproductive experience' of women who work is better than those who do not.⁵⁸ Some studies show that women who are employed have a lower risk of preterm delivery than those who are not.^{23,38,138} However, a pregnant woman may be exposed, while at work, to particular hazards that might potentially cause adverse outcomes for mother or fetus. Therefore, where possible, steps should be taken to minimise exposure where there is sufficient evidence that the risk of maternal or fetal harm outweighs any benefit to health. A few workplace exposures (eg lead and ionising radiation)^{1,2} are well established as being harmful during pregnancy and the need to limit exposure is explicit in statutory instruments in the UK. However for many other common workplace exposures, including some physical hazards, the evidence to inform a balanced assessment of risk is not available, or is uncertain.

Legal framework

A number of statutory instruments in the UK relate directly to the management of pregnant women at work. Employers have a statutory obligation, as governed by the Management of Health and Safety at Work Regulations 1999,³ to protect the health and safety of new and expectant mothers. Once informed in writing, the employer is obliged to undertake an

individual risk assessment that aims to identify hazards and assess risks. The employer must take action to remove, reduce, or control any risks that are significant. In the event that significant risks cannot be controlled adequately, the employer has a duty to undertake a hierarchy of measures comprising temporary adjustment in working conditions, provision of alternative work or suspension on paid leave. The Employment Rights Act 1996⁴ (amended by the Employment Relations Act 1999)⁵ requires employers, when offering suitable alternative work, to ensure that this is appropriate and offered on terms and conditions that are no less favourable than the new or expectant mother's normal terms and conditions.

The Workplace (Health, Safety and Welfare) Regulations 1992⁶ require employers to provide suitable rest facilities for workers who are pregnant or breastfeeding. These must be suitably located (eg near to toilets) and, where necessary, should provide appropriate facilities for the new or expectant mother to lie down.

Under the Sex Discrimination Act 1975⁷ any breach of health and safety legislation in relation to new and expectant mothers is considered automatically to be sex discrimination.

The Council Directive 92/85/EEC of 19 October 1992 states that, '...pregnant workers [and] workers who have recently given birth or who are breastfeeding must be considered a specific risk group...and measures must be taken with regard to their safety and health.' It further advises that a risk assessment must be undertaken and that the worker is protected.⁴³

Existing guidance

The Health and Safety Executive (HSE) provides guidance on risk assessment in respect of pregnant, newly delivered and breast-feeding mothers. This is provided in the form of two booklets which are targeted at employees and health advisers respectively: *A guide for new and expectant mothers who work*,⁷¹ and *New and expectant mothers at work – a guide for health professionals*.⁷²

However, with the exception of hazards that are governed by specific legislation, there are no clear guidelines regarding the level of risk at which adjustments to work should be made for pregnant workers, or what adverse outcomes might result from poor risk control. Moreover, existing guidance documents do not describe the scientific justification for decisions about risk controls. There is, therefore, a need for new evidence-based guidelines in this area. These will help to inform both employees and employers more fully of the nature and level of risk to pregnant and newly delivering mothers who work, and their offspring.

Epidemiology of adverse pregnancy outcomes

Fetal outcomes

The adverse outcomes that are measurable immediately after the end of a pregnancy (and which feature in the research literature on workplace hazards) include spontaneous miscarriage, preterm delivery, still birth and low birth weight. Approximately 15% of medically recognised pregnancies end in spontaneous miscarriage,⁷⁴ and around 7% of babies are delivered preterm. Figures from 2000 indicate that this rate is lower in Finland (6.3%), Sweden (5.8%), Denmark (3.3%), and Iceland (3.6%) than in the UK (7%).⁶⁰ In 1998, in England and Wales, 7.48% of live births were of low birth weight (<2500 g)⁹⁰ and the rate of still births per 1,000 total births was 3.9.^{20,90} The UK has the highest rate of preterm delivery in Europe (www.tommys.org).

These adverse birth outcomes are clinically important. They are recognised risk factors for poor health in the perinatal period, childhood and even later in life. Birth weight⁸⁶ and gestational age are the most important neonatal risk factors that determine perinatal mortality.^{83,169} Preterm birth is a major contributor to perinatal mortality and morbidity.^{60,65,98,108,133} Low birth weight is related to neonatal mortality, a significant determinant of post-neonatal mortality and infant and childhood morbidity.¹⁶¹ Moreover, there is considerable evidence that low birth weight predicts poor growth and development, and increases the risk of chronic diseases in adulthood, including coronary heart disease and stroke, hypertension, non-insulin-dependent diabetes, obstructive lung disease and neurological and cognitive impairment.^{44,163}

Maternal outcomes

A number of complications of pregnancy that might occur during the period that women are still working can be regarded as adverse maternal outcomes. These include gestational hypertension and pre-eclampsia (gestational hypertension with proteinuria and oedema). Preterm delivery and spontaneous preterm prelabour rupture of membranes may pose a threat to both fetal and maternal health, for example by increasing the risk of intra-uterine infection, which may in turn lead to maternal and fetal sepsis.

Other outcomes

Complications of pregnancy may lead to an early or prolonged period of sickness absence or delayed return to work. These could be regarded as an adverse outcome for both the employee and employer. The tendency to take absence during pregnancy is strongly influenced by health, beliefs and health advice. Therefore, any relationship between workplace hazards and excess absence from work is likely to be biased by individual and societal perceptions of risks to a pregnancy.

General risk factors

There are a number of well-established predictors of adverse pregnancy outcome. These include maternal age, lower education level, low and high body mass index (BMI), smoking and low socio-economic status.^{86,88} Others are specifically related to preterm birth⁶⁰ or to low birth weight, namely race, father's occupation, education level, history of previous miscarriage or preterm birth and, possibly, age and parity,¹⁰⁸ cigarette smoking, short maternal stature and poor nutrition before and during pregnancy.⁸⁶ Some of these factors are clearly more amenable to intervention than others, for example smoking. However, in theory, the majority of hazards to pregnancy that are encountered in the workplace can be modified.¹⁴⁰

Occupation and pregnancy

A number of studies have pointed to an increased risk of adverse pregnancy outcomes in certain occupations. The McDonald series of papers^{103–108} noted that, for fetal death, the odds ratios were significantly increased for nursing aides and orderlies, food and beverage servers and workers in factories ($p < 0.01$). Late, but not early, spontaneous miscarriages were significantly raised in radiology technicians ($p < 0.01$) and operating room nurses, and those working in agriculture and horticulture ($p < 0.05$).¹⁰⁷

For preterm birth, the observed/expected (O/E) ratios were significantly raised in the food and beverage services ($p=0.03$) and psychiatric nurses ($p<0.01$). The O/E ratios for low birth weight were also significantly raised in the food and beverage services ($p<0.05$), chambermaids and cleaners ($p=0.03$) and metal and electrical manufacturers ($p<0.01$).¹⁰⁷

Nurses (excluding psychiatric and operating room nurses) and childcare workers had significantly lower mean percentage predicted birth weight for age ($p<0.05$).¹⁴

With regard to spontaneous miscarriage, odds ratios were significantly raised for both previous and current pregnancies in nursing assistants and attendants, and food and beverage servers ($p<0.05$).¹⁰⁴

Manual workers and healthcare workers have been found to have a significantly higher risk of intra-uterine growth restriction.¹⁵²

Possible biological mechanisms

It is not entirely understood how the workplace exposures considered in this guideline could result in adverse pregnancy outcomes. A number of biological/physiological mechanisms have been hypothesised, although there is little direct evidence to support them. Plausible explanations for and against a causal pathway for the adverse outcomes of interest are summarised below.

Prolonged standing

It is well established that prolonged standing results in pooling of blood in the veins of the legs, potentially resulting in reduced venous return, cardiac output, and ultimately arterial blood pressure. This effect might be potentiated by the presence of a gravid uterus, particularly in the third trimester of pregnancy. It has been postulated that, under some circumstances, this could lead to a reduction in uterine blood flow. In theory this might increase uterine contractility (and therefore the risk of spontaneous miscarriage or preterm delivery), or potentially reduce the rate of fetal growth.⁵⁸ However, it could be argued that strong physiological mechanisms exist to protect fetal blood flow in pregnancy, and that the extent and duration of maternal hypotension required to override this adaptation would be extreme. It is not clear whether this proposed mechanism is likely to occur in the context of prolonged standing at work under normal environmental working conditions.

Hard physical work

Altered body posture and heavy physical exercise or strain might reduce maternal blood pressure and/or blood flow from the uterus to the placenta. It is possible that resulting under-perfusion could result in restricted fetal growth and impaired survival.^{9,36,42,68,86,115} There may also be increased substrate utilisation by muscles⁶⁸ with increased maternal energy requirements.⁸⁶ Theoretically this might use up calories needed by the fetus,¹¹⁵ resulting in nutritional deficits.⁹ These physiological changes are potentially intensified either by an increase in workload or duration of physical activity. However (as for prolonged standing), in order to be a significant threat to fetal growth these changes would have to override the normal physiological adaptations to pregnancy that aim to preserve fetal oxygen consumption and nutrition.

It has been postulated that physical strain might also cause spontaneous miscarriage by means of increased abdominal pressure, elevated body temperature or sudden change in hormones (increased norepinephrine levels).⁶⁸ A further possible mechanism is that physical strain may precipitate uterine contractions, and hence premature cervical effacement and dilatation.¹⁵⁰

Heavy lifting

It is known that heavy lifting causes a temporary increase in intra-abdominal pressure when the subject holds their breath against a closed glottis (Valsalva manoeuvre). Although there is no direct evidence, it has been postulated that raised intra-abdominal pressure might in turn provoke uterine contractions.⁹

Shift work

It has been postulated that shift work might influence reproductive function in humans through hormonal disturbances.^{15,25,74,171,172} Both direct (through changes in circadian rhythm) and indirect (through psychosocial stresses and disturbed sleep)¹¹⁹ mechanisms have been proposed. Zhu *et al* suggested that an increase in oestrogen levels may be linked to adverse pregnancy outcomes.¹⁷¹ The authors postulated the following causal mechanism: suppression of nocturnal melatonin production by light during the night could lead to increased oestrogen levels which stimulate the growth of hormone-sensitive tumours. Since oestrogen supplementation during pregnancy has been shown to increase the risk of spontaneous miscarriage, increased levels of oestrogens due to night work theoretically could increase the risk of fetal loss.¹⁷¹

In summary, a number of mechanisms have been proposed that might explain a causal relationship between some physical aspects of work, or shift work, and adverse pregnancy outcomes. However, these would involve overriding of the normal physiological adaptations to preserve the fetus. Overall it is uncertain whether this would be likely to occur in a work situation. Indeed it seems relatively unlikely unless physical exposures and environmental conditions were particularly severe.

Evidence of variation in practice: relevance to occupational medicine

There is some evidence of inconsistency in the advice given to pregnant women, dependent upon who undertakes the risk assessment and how it is performed. For example, in a study by Conrad,³⁸ a questionnaire was sent to lead clinicians in NHS occupational health departments in the UK. The questionnaire enquired about the presence of written policies regarding the management of pregnant healthcare workers, whether alternative work was available to replace normal duties, who was responsible for risk assessments and whether work restrictions were suggested with regard to particular hazards. The results indicated that there was substantial variation in practice across the NHS.

In another study in the USA in 1998, a survey was mailed to physicians who were part of either family residency programmes or obstetrics-gynaecology programmes. The physician was given a randomly selected case study and asked whether or not they would recommend restrictions. They were more likely to recommend job restrictions if there were risk factors for preterm birth. However, recommendations about restrictions were not consistent. Variation was most pronounced if there was low obstetrical risk, presumably because recommendation then

Physical and shift work in pregnancy: occupational aspects of management

depended upon the physician's own perception of risk. For example among physicians who said they would always recommend restrictions for low risk, 35.4% believed that standing is an important risk factor, compared with 6.7% of physicians who said they would sometimes, rarely or never restrict. There were similar beliefs regarding lifting.⁵² Evidence of variation of practice is also found in a Danish study.¹⁶⁰

One method for improving the consistency of practice is to use evidence-based guidance as a baseline for a continuing cycle of audit and review. This approach has been shown to improve clinical effectiveness in other specialties.¹³⁴ The same approach will be adopted in relation to occupational health advice for pregnant women who are exposed to hazards at work, using this guideline as a starting point.

2 Methodology of the evidence review

The population under study was defined as pregnant women at work.

In order to define which hazards and outcomes would be addressed, key questions were formulated by the GDG Leader. After discussion amongst GDG members, a list of possible adverse outcomes was considered. By consensus, a number of outcomes including caesarean section, instrumental delivery, psychological symptoms and stress were omitted. An initial literature search was undertaken to assess the potential volume of available literature for the remaining questions and outcomes of interest. Following this scoping search, the list of questions was revised and agreed by the GDG. The final key questions are shown below.

The key questions narrow the scope of the guidelines to a few hazards only, but the approach could be extended to incorporate other hazards if the guidelines were to be developed further in the future. Similarly the list of adverse outcomes could be expanded in future revisions of the guidelines.

For each of these hazards, the evidence base was assessed to explore the level of risk for each adverse pregnancy outcome. The volume, strength and consistency of the evidence was considered with a view to suggesting recommendations for appropriate adjustments in the workplace.

Key questions

1. What is the effect on mother and fetus of manual handling and prolonged standing?

Population	Pregnant women at work
Hazards	Manual handling Prolonged standing
Outcome	<ol style="list-style-type: none">Fetal morbidity<ol style="list-style-type: none">spontaneous miscarriagelow birth weightIUGR/SGA*preterm birthstillbirthMaternal morbidity<ol style="list-style-type: none">gestational hypertension/pre-eclampsiapreterm delivery to include threatened preterm labour and preterm prelabour rupture of membranesSickness absence

*IUGR = intra-uterine growth retardation; *SGA = small for gestational age. SGA fetuses are small but not necessarily growth restricted, whereas IUGR is defined by serial ultrasound measurements showing diminishing fetal growth.¹⁵²

2. What is the effect on mother and fetus of the mother working more than eight hours on a shift, more than 37 hours a week, or shift work (to include any shift pattern).

Population	Pregnant women at work
Hazards	Working more than eight hours on a shift Working more than 37 hours a week Working any shift pattern
Outcome	<ol style="list-style-type: none"> 1. Fetal morbidity <ol style="list-style-type: none"> (a) spontaneous miscarriage (b) low birth weight (c) IUGR/SGA (d) preterm birth (e) stillbirth 2. Maternal morbidity <ol style="list-style-type: none"> (a) gestational hypertension/pre-eclampsia (b) preterm delivery to include threatened preterm labour and preterm prelabour rupture of membranes 3. Sickness absence

3. What is the effect on mother and fetus of the mother working night shifts?

Population	Pregnant women at work
Hazard	Working night shifts
Outcome	<ol style="list-style-type: none"> 1. Fetal morbidity <ol style="list-style-type: none"> (a) spontaneous miscarriage (b) low birth weight (c) IUGR/SGA (d) preterm birth (e) stillbirth 2. Maternal morbidity <ol style="list-style-type: none"> (a) gestational hypertension/pre-eclampsia (b) preterm delivery to include threatened preterm labour and preterm prelabour rupture of membranes 3. Sickness absence

The GDG agreed upon standard definitions for maternal and fetal outcomes using standard texts,^{90,30} and sought the opinion of the obstetric consultant on the GDG if any definitions were unclear (see Glossary).

Literature search

To form the evidence base, a systematic literature search was undertaken. The terms in the key questions were used as the basis of the literature search.

The search terms and the databases to be searched were defined in collaboration with a Department of Health (DH) librarian. An initial search was carried out in 2004. A repeat search was undertaken the following year (2005) by another librarian to update the evidence base. In addition the GDG was made aware of a high quality meta-analysis²⁷ that had been submitted for publication in 2006 (published 2007) which it was decided should be included in the

evidence base. The terms were searched as both text and Medical Subject Headings terms (MeSH)[®], www.nlm.nih.gov). The numbered list was searched with all the other terms/phrases in the subsequent list. (See Appendix 1 for complete search strategy and databases searched.) A comprehensive evidence-based policy on the management of expectant women from John Lewis Partnership was hand searched. Other papers identified from this search were included if relevant. The following sources were searched for grey literature:

- (1) Royal Free Hampstead NHS Trust Research and Development database
- (2) Faculty of Occupational Medicine Database of Membership of the Faculty of Occupational Medicine (MFOM) dissertations
- (3) Institute of Occupational Safety and Health (IOSH) website
- (4) Specifically searching for articles authored by Linda Tapp (manual handling and ergonomics expert as advised by the GDG)
- (5) General enquiries made to other colleagues.

The inclusion/exclusion criteria for papers are shown in Table 1.


Table 1 Inclusion/exclusion criteria for papers

Inclusion criteria	Exclusion criteria
1980–2006*	Does not answer key questions
Observational studies, intervention studies and literature reviews	Multiple pregnancies
Human only	Data collected prior to 1980
Randomised controlled trials (RCTs), case-control studies, longitudinal studies, meta-analyses, review articles	History of chronic illness
	Women not employed during pregnancy
* A high quality meta-analysis that had been submitted for publication in 2006 was included in the evidence base for the guideline.	

Following the literature searches, 270 abstracts were found to be relevant to the key questions and fitted the above inclusion criteria. All of the abstracts were read by two GDG members, Paul Grime (PG) and Nadia Sheikh (NS), to reduce bias in literature collection and to ensure consistency in selection of the papers for full appraisal. Having identified papers relevant to the key questions, all were read by the GDG Leader (NS) to determine whether they should be included as part of the critical appraisal. Papers that were not considered to be relevant to the key questions were excluded, and the remainder were critically appraised (see Fig 1). Although some papers were not included as evidence, they still contained some useful/interesting material, and have therefore been included in the introduction.

One paper was included as evidence, even though data was collected prior to 1980, because it was referenced in a number of credible papers and also formed the basis of subsequent research by others. The consensus opinion of the GDG was that in order to help understand the evidence in subsequent papers using the same methodology, this paper should be appraised and included as evidence.⁹⁸ It is recognised that the inclusion of this paper may introduce some selection

Fig 1 Flow chart for study selection



bias, however as there is an extensive body of papers involved in the review the GDG considered that the inclusion of this paper would not importantly affect the overall conclusions of the literature review.

Foreign language articles were included in the initial literature search; however these were excluded prior to the critical appraisal stage due to the variety of languages, resource and time limitations for translation.

Due to the nature of the key questions that were being considered, no randomised controlled trials were identified. This is not uncommon in the literature relating to occupational hazards, most of which comprises observational studies. The majority of papers identified in this literature search were case-control or cohort studies. One meta-analysis was identified, but this was excluded after critical appraisal. One systematic review, which has been included, also includes a meta-analysis of some of the exposure-outcome combinations. The majority of review articles that were identified did not follow systematic methodology, and therefore could not be included as evidence. However, relevant references cited in these narrative reviews were retrieved and appraised accordingly.

The number of excluded papers is shown in Table 2.

Table 2 Excluded papers

Does not address key questions	Initial reason for exclusion						TOTAL EXCLUDED
	Appraised: 2- or 1- indicating a high risk of bias or confounding	Study population not pregnant women at work	Not original research or not systematic review/ editorial/ book review	Foreign language	British Library unable to retrieve paper	Very old data	
44	25	3	30	8	2	1	113

Critical appraisal

Each included paper was critically appraised using a standard form (see Appendix 2). Each paper was appraised independently by two committee members, who then discussed the paper and agreed upon a final grade. A total of eight people were involved, to varying degrees, in appraising the evidence. All appraisers received training on critical appraisal techniques. The majority (approximately 75%) of the papers were appraised by Nadia Sheikh (NS) and Paul Grime (PG). NS also reviewed the papers that were appraised by other GDG members, to ensure consistency and accuracy.

Assessing the evidence

Various methods for assessing and grading the evidence in papers have been published. These include grading methods produced by:

- The Health Evidence Network
- Human Fertilisation and Embryology Authority (HFEA) executive
- Scottish Intercollegiate Guidelines Network (SIGN)
- US Government Agency
- NHS Research and Development Centre

All describe similar methods of grading papers based upon the type of paper, eg RCT/meta-analysis, and the quality of the study. The SIGN methodology was used to assess the evidence for this guideline, primarily because it is suitable for use when appraisal will be carried out by members of the GDG (rather than a dedicated academic team). All GDG members who participated in critical appraisal received standard training in critical appraisal techniques. The SIGN grading system is reproduced in Appendix 3.

One of the concerns about the SIGN methodology is the relative lack of transparency of the simple quality scoring system, which does not state in detail the rules for assessment of major methodological issues such as bias and confounding. Therefore, in addition to the SIGN rating, a scoring system was devised for use in assessing the evidence for this guideline. The purpose of this system was to demonstrate more clearly the consistent application of the SIGN quality rating. This additional scoring system was applied to all of the papers that were included as evidence. The detailed scoring tables can be found in Appendix 5.

Development of recommendations

Draft recommendations were produced by the GDG Leader and then circulated to the GDG members for comments and feedback. The draft guideline document was reviewed by three external assessors and the steering group of the Occupational Health Clinical Effectiveness Unit (OHCEU), and revised in response to their comments.

Limitations of the evidence base

General

Heterogeneity

There was considerable heterogeneity among the body of studies that was included in the critical appraisal and final evidence tables. Populations varied in respect of country and occupational group. The range of exposures within a particular type of job might vary by country, thus affecting generalisability to a UK population. However, this was not a major problem as relatively few studies used job title as the main method of exposure assessment.

Outcome definitions and method of outcome assessment also varied, for example both ultrasound scanning and the Dubowitz score were used in different studies as measures of fetal growth (the vast majority of studies used ultrasound for measurement \pm last menstrual period, only a few of the older studies used a Dubowitz score). Exposure categories and methods of exposure assessment also varied. This reduced the potential for combining findings in a formal meta-analysis, although one high quality study that did undertake a meta-analysis of a subset of the papers with similar outcome and exposure measures was included. The heterogeneity of outcome and exposure assessment might explain some of the inconsistency between the results of studies that is described in the summary of the findings from the evidence review.

Timing of exposure in relation to different stages of pregnancy

Workplace hazards may present risks at different stages of pregnancy. For example, there may be a greater risk in the early stages of pregnancy when the fetus is not yet fully formed. If a study looks at a particular point of time during pregnancy, for example in a cross-sectional survey, it may miss an association that is present at either an earlier or a later stage. Many of the studies in this review did not include detailed information about the timing of exposure in relation to pregnancy. Where this information was given it has been included in the evidence tables. In addition, the failure to note the timing of the exposure makes it difficult to advise on the application of workplace restrictions/controls.

Bias

A common problem in this literature was the reliance on subjective recall for the assessment of exposure after the outcome of the pregnancy was known (typically by questionnaire at the time of delivery or later). This introduced an important potential for inflationary bias. Mothers who had an adverse outcome might be expected to recall exposures more clearly, or even over-estimate their previous exposures, if they perceived a harmful effect of the exposure on their pregnancy. This could lead to over-estimation of the association between the adverse outcome and the exposure of interest.

Confounding

One of the most striking aspects of the review is the number of potential confounding factors. These include: extremes of maternal age (<20 and >35), lower education level, low pre-pregnancy BMI, smoking, alcohol consumption and low socio-economic status.

Not all papers addressed all of the potentially important confounding factors. Confounders are likely to be most important if the strength of their associations with the risk factor of primary interest, and with the health outcomes, are strong and if the prevalence of the confounder in the population of interest (in this instance working pregnant women) is high. If a study design did not adjust for potential confounders the GDG looked to see if a reasonable justification was provided. For example, in a Chinese study smoking was not thought to be an important confounder because very few women smoked.¹⁷⁰ Failure to attempt to identify or address confounders, or insufficient justification for failure to address potential confounders, was reflected in the grade awarded to the paper. If the appraisers judged there to be a reasonable likelihood that results from a given study were strongly influenced by confounding, the paper was given a low score (1– or 2–) and was excluded from the final evidence tables.

Parity

Some adverse outcomes may be more likely in second and subsequent pregnancies, for example the increased risk of miscarriage if there is previous history. Women may already have modified their job by subsequent pregnancies in light of hazards encountered in the previous pregnancy. Pregnant women may choose to request adjustments at an earlier or a later stage of pregnancy, depending upon the outcome of previous pregnancies. Therefore parity needs to be adjusted for, or at the very least acknowledged as a potential confounder. This was done in most cases, so parity is unlikely to have had an important influence on these results.

Healthy worker effect

It is possible that a significant association was not found in some cases because those women still working during pregnancy are doing so because they are 'healthy' and therefore less likely to experience adverse outcomes. This would tend to lead to an under-estimate of risk.

Exposure assessment

Misclassification bias

Exposure misclassification is more likely to occur with certain methods of exposure assessment, for example using job title alone to assess complicated exposures. As only one paper included evidence assessed on the basis of job category this is not thought to be an important factor in this review.

Night shifts

When studying night shifts, most papers examined the shift pattern rather than the intensity of work being undertaken. Most studies expected the night shift to have the most significant effect, but a few noted that the evening shift had the worst associated outcomes. This could be a reflection of the intensity of work undertaken on that particular shift. Those papers where a significant association was noted were conducted on hospital employees.

Where a woman was working less than 40 hours per week, the out-of-hours work per day was not necessarily quantified.

Physical effort

This was not always quantified in detail, and was generally just classified as light, moderate or heavy. Some papers defined these categories; others did not. One paper defined moderate physical exertion as lifting or carrying anything equal to or more than 10 kg at least three times a day.¹³³ Another measured physical intensity on the type of physical effort: carrying heavy loads, performing heavy cleaning tasks and sweating whilst undertaking physical activities.¹⁵²

Complexity of exposure assessment

The above comments in relation to night work and physical effort are examples of how complex exposures, consisting of more than one aspect (eg intensity and duration) are oversimplified in assessment for research purposes. This problem also relates to some of the other exposures in this review. If the risk of adverse pregnancy outcome is specific to certain aspects of exposure, and these aspects are not differentiated between as a result of crude exposure assessment, then important effects might be missed due to dilution.

Outcome

Difference between small for gestational age (SGA) and intra-uterine growth restriction (IUGR)

Studies tend to consider one or the other of these as adverse outcomes, even though they are not necessarily the same outcome. SGA fetuses are small but not necessarily growth restricted. Therefore serial ultrasound measurements showing diminishing fetal growth, compared to SGA, may better reflect the occurrence of restricted growth, ie IUGR.¹⁵²

3 Findings and recommendations

Synthesis of the evidence and recommendations

In making recommendations for each of the exposures of interest, a number of factors were taken into account. The quality and consistency of the evidence for similar exposures and outcomes were considered. High quality studies and large studies (high number of pregnancies observed) were given more weight than lower quality or smaller investigations. Statistical significance was not considered in isolation; the magnitude of the associations (size of the risk estimates) and the likely clinical importance influenced the synthesis of evidence strongly. The GDG was cautious not to give too much weight to risk estimates that were rather close to unity (1.3 or less), as it is recognised that associations of this order in observational studies are often due to unrecognised bias or confounding.

In deciding how to approach the uncertainties in the evidence base, the GDG took into account the fact that many of the exposures of interest (for example lifting) are common in the normal daily life of mothers, particularly if they already have young children at home. It would be inappropriate to alarm women about normal everyday activities unless the evidence of harm is compelling. Moreover, it is possible that remaining active during pregnancy, even if it involves, for example, prolonged standing or lifting, may actually be beneficial. Clearly a high level of certainty is required before recommendations are made to exclude pregnant women from activities if these restrictions might actually be bad for health.

1 Manual handling, lifting, heavy physical work

This review identified a total of 17 original papers and one systematic review that addressed the association between manual handling, lifting or heavy physical work and the outcomes of interest. The high quality systematic review²⁷ included six of the 17 papers identified by this review and 28 additional papers, giving 45 in total that addressed lifting/physical activity in relation to adverse birth outcomes.

(a) Heavy physical effort

Four studies^{83,107,133,141} were found (three 2+ and one 2++) that explored the association between physical activity and preterm birth/preterm labour or spontaneous miscarriage (Evidence tables 2 and 3). Of these, three showed a significant positive association,^{83,107,133} including risk estimates of 2.91 and 1.87. However there was a substantial body of conflicting evidence that showed no association between heavy physical work and preterm birth, or suggested only a small increase in risk. One paper¹⁴¹ (2+) identified by this review did not find a significant increase in preterm births according to heavy working conditions. A high quality systematic review²⁷ (1++) found 20 additional papers (excluding one of the studies in Evidence table 2) that addressed the link between physical workload and preterm delivery. The authors' tables demonstrated that 12 of the studies found risk estimates close to unity (<1.3), and the remainder (eight studies) showed a positive association with risk estimates >1.5. The authors pointed out that risk estimates tended to be lower (<1.3) in prospective studies, and that four of the five largest studies also had rather low risk estimates (<1.3).²⁷ They concluded that the evidence was less consistent than for other exposures and that more research would be helpful.

Three studies^{86,138,152} (2+) (Evidence tables 4 and 6) examined the association of physical activity with low birth weight/IUGR or SGA. All three showed a positive association, with an odds ratio of 2.4 in one. A high quality systematic review²⁷ (1++) identified an additional seven studies (not included in Evidence table 6) that addressed the association of physical activity with SGA. Three of these studies were positive, showing significantly increased risk between 1.32 and 2.4. In contrast, four studies were negative, with relative risk less than unity.

The same systematic review²⁷ (1++) identified five studies that explored the relationship between physical activity and pre-eclampsia or pregnancy-induced hypertension. Of these, two showed a significant positive relationship, with relative risks 2.1 and 3.47 respectively. However, three studies were negative.

The same systematic review judged that exposure assessment for heavy physical work was too heterogeneous to allow meta-analysis.²⁷

Conclusion

There was a large body of evidence (24 papers including those identified within a systematic review) that addressed the association of heavy physical work with preterm delivery. However, there were some conflicting findings. Overall rather more studies did not find a significant association (13/24) with this outcome than were positive (11/24). Among the studies that were positive, risk estimates were generally ≤ 2 . There were fewer papers (10) about low birth weight, IUGR or SGA and again there was some inconsistency, although rather more papers showed a positive association (6/10) than were negative (4/10). Overall the positive papers pointed to a moderate effect (relative risks up to 2.4). For pre-eclampsia and gestational hypertension there were a limited number of papers (five in all, all identified within a systematic review) and the evidence was inconsistent (2/5 positive and 3/5 negative).

RECOMMENDATIONS

Recommendation	Risk	Grade	Evidence
Pregnant workers should be informed that evidence suggests that heavy physical work carries no more than a moderate risk of low birth weight/IUGR/SGA, but there is only limited and inconsistent evidence of risk for preterm birth and pre-eclampsia	Large body of evidence with some conflicting findings, but reasonably consistent estimates of risk that tend to rule out more than a moderate risk of low birth weight/IUGR/SGA.	B	152, 138, 86, review ref 27 (includes a further 7 papers)
Employers should reduce very heavy physical activities for pregnant workers where possible, particularly in late pregnancy. However, if a pregnant worker who has been informed of the possible risks wishes to continue then there are insufficient grounds upon which to impose restrictions against her will	Large body of studies, but inconsistent findings in relation to risk of preterm delivery. Limited number of studies and inconsistent evidence of risk of pre-eclampsia or pregnancy-induced hypertension.		83, 107, 141, 133, review ref 27 (includes a further 20 papers) Review ref 27 includes 5 papers

The exposure measures for heavy physical work are too heterogeneous to inform a recommendation specifying a level of physical work at which adjustments to work should be implemented.

(b) Lifting

Three studies^{9,108,161} were found (one 2++ and two 2+) (Evidence table 2) that explored the relationship between lifting and preterm birth or labour. Of these, two showed no significant association^{9,161} and one showed a significant association with a small increase in risk (O/E 1.26).¹⁰⁸ In addition, four studies explored lifting and manual handling and spontaneous miscarriage/perinatal mortality (Evidence table 3). Of these, three showed positive associations^{104,107,170} (2+) with risk estimates between 1.5 and 2.0 and one showed no evidence of a relation with frequently lifting >15 lbs⁴² (2+). A high quality systematic review (1++) found a further seven studies that examined lifting in relation to preterm delivery.²⁷ The authors tabulated results from these studies, showing risk estimates below 1.15 in six studies and between 1.3 and 1.49 in two studies. In both of the latter, 95% confidence intervals included 1, so the findings were not statistically significant.

Four studies addressed the association of lifting with low birth weight/IUGR/SGA (Evidence tables 4 and 6). Of these, only two found important positive associations^{14,161} (2+) with relative risk 3.5 (risk estimates were not given for the second paper), and a third⁹ (2+) was inconclusive. A fourth paper¹⁰⁸ (2++) found a very small increased risk (relative risk 1.25). A high quality systematic review²⁷ (1++) identified a further three studies (excluding those in Evidence tables 4 and 6) that examined lifting and SGA: all three papers found no significant association.

One paper (Evidence table 5) explored lifting in relation to pre-eclampsia¹⁶⁵ (2+). This study found a positive association with lifting heavy loads in early pregnancy (odds ratio 1.7) and with lifting heavy loads >20 times weekly in early pregnancy (odds ratio 2.0). A high quality systematic review²⁷ (1++) identified one further study (in addition to the study in Evidence table 5) that examined lifting and pre-eclampsia/pregnancy-induced hypertension. This study showed no association with pre-eclampsia and a significant negative association with pregnancy-induced hypertension (relative risk 0.68).

A systematic review²⁷ considered that lifting outcomes were too heterogeneous to enable formal meta-analysis.

This literature search identified three studies that looked at the association between lifting or manual handling and sick leave/sick certification.^{77,138,155} Two studies showed that pregnant women who lifted heavy loads at work were more likely to take time off sick (odds ratio 1.9)⁷⁷ (2+), or to leave work >3 weeks before delivery (odds ratio 1.48)¹⁵⁵ (2+). One of these studies¹⁵⁵ found that if work involved manual handling activities (twisting and bending or working with the hands above shoulders), pregnant women were more likely to leave work >3 weeks before delivery (odds ratios 1.46 and 1.55 respectively) and >8 weeks before delivery (odds ratios 1.32 and 1.36 respectively). This finding does not provide direct evidence of actual harm to the mother or fetus, and probably reflects the societal beliefs about the effects of heavy work on pregnancy. It is not therefore useful in forming recommendations about restriction from work on medical grounds during pregnancy.

Conclusion

There was a substantial body of evidence for preterm delivery, with general consistency of results among 14 studies identified. These showed either no statistically significant association with lifting (9/14) or a modestly elevated risk (6/14, although of these, two were not statistically significant). The evidence in relation to SGA was limited to seven studies. These were generally consistent, showing a weak positive association (3/7), no association (3/7) or inconclusive (1/7). There were very few papers (two papers) addressing pre-eclampsia and pregnancy-induced hypertension, and these gave conflicting results (one positive and one negative association).

RECOMMENDATIONS

Recommendation	Risk	Grade	Evidence
Pregnant employees should be informed about the generally consistent evidence suggesting that lifting carries no more than a moderate risk of preterm birth and low birth weight, but limited inconsistent evidence for pre-eclampsia	Large and generally consistent body of evidence rules out more than a modest risk (relative risk <1.3) of preterm delivery	B	9,161, 108, 104, 107, 170, 42 review ref 27 (includes a further 7 papers)
Employers should reduce lifting for pregnant workers where possible, particularly in late pregnancy. However, if a pregnant worker who has been informed of the possible risks wishes to continue then there are insufficient grounds upon which to impose restrictions against her will	Reasonable body of evidence, generally consistent in suggesting no effect for low birth weight/IUGR/SGA. Very limited evidence for pre-eclampsia/pregnancy-induced hypertension		14, 108, 161, 9, review ref 27 includes a further 3 papers 165, review ref 27 includes 1 further paper

The exposure measures for lifting work are too heterogeneous to inform a recommendation specifying the actual level of physical work at which adjustment to work should be implemented.

2 Prolonged standing

One systematic review and 11 original papers were identified that examined the relationship between prolonged standing or standing and walking and the adverse pregnancy outcomes of interest.

Five papers addressed preterm delivery^{65,98,133,140,159} (three 2+ and two 2++) (Evidence table 2) and five papers addressed spontaneous miscarriage/perinatal mortality^{11,42,46,104,107} (four 2+ and one 2++) (Evidence table 3). Prolonged standing was defined variously as >3 hours (3 papers), >5 hours (1 paper) or >6 hours (1 paper). All five studies for preterm delivery showed a statistically significant positive association between preterm delivery and risk estimates 1.26–4.10 (two below 2.0 and three above 2.0). All five studies for spontaneous miscarriage/perinatal mortality showed a statistically significant association with risk estimates 1.32–4.32 (two below 1.5 and three above 1.5). A high quality systematic review²⁷ (1++) identified a further 16 papers (in addition to those in Evidence tables 2 and 3). The authors describe moderately elevated risks of preterm delivery in seven studies with relative risks >1.5. However, they comment that in the largest studies relative risks were in the range 1.07–1.56. They also carried out a meta-analysis

across 12 studies that compared standing for at least three hours, and found a pooled estimate of risk of 1.28 (95% CI 1.11 to 1.47; $Q=16.50$, $p=0.12$).

Three studies addressed low birth weight^{58,65,159} (2+) (Evidence table 4). Two showed a significantly increased risk with standing. The risk estimate was not stated in one study⁵⁸, but in the other was 3.3.⁶⁵ The third¹⁵⁹ showed a positive association between standing and low birth weight (relative risk 1.40; 95% CI 0.64 to 3.03). The association in the standing group compared with the sedentary group was not statistically significant at the 5% level. A high quality systematic review²⁷ (1++) identified a further eight papers (that were not identified in this review) relating to SGA, including four of high quality. Of these, six showed no effect with relative risks close to unity. Two studies had statistically significant relative risks, one of 1.4, and one a relative risk of 2 that was not statistically significant.

No original papers were found for pre-eclampsia, but this outcome was addressed in a high quality systematic review²⁷ (1++). The authors found five studies, although only one was of high quality. None of the studies showed a significant positive risk from prolonged standing.

Conclusion

There was an extensive body of evidence for the association of standing with preterm delivery, spontaneous miscarriage and perinatal mortality for which the findings were broadly consistent. The evidence pointed to no more than a moderate risk from prolonged standing (14/21 studies, risk estimates <1.5). Moreover it tended to rule out a large effect (high quality meta-analysis with pooled risk 1.28). A reasonable body of evidence for low birth weight was also consistent, and suggested no effect (6/11 studies negative and at least one further study with risk estimate <1.5). A small body of evidence for pre-eclampsia was limited in quality (although included one high quality study) but showed no effect.

RECOMMENDATIONS

Recommendation	Risk	Grade	Evidence
Pregnant employees should be informed about the generally consistent evidence suggesting that prolonged standing (>3 hours) carries no more than a small risk of preterm birth and low birth weight/ IUGR/SGA, and limited evidence for no effect for pre-eclampsia	Good evidence suggesting no more than a small risk of preterm birth	A	11, 42, 46, 65, 98, 104, 107, 133, 140, 159, review ref 27 includes a further 16 papers
If employees stand for >3 hours/day, employers should consider reducing this or providing alternative work for that period of time, to reduce hours of standing to the minimum possible	Large body of consistent evidence suggesting no risk or no more than a small risk of low birth weight		58, 65, 159, review ref 27 includes a further 8 papers
However, if a pregnant worker who has been informed of the possible risks wishes to continue then there are insufficient grounds upon which to impose restrictions against her will	Limited evidence suggesting no effect for pre-eclampsia		Review ref 27 includes 5 papers

From current evidence it cannot be stated conclusively at which stage of pregnancy these recommendations should be implemented. Most studies used three hours or more to define prolonged standing, so it seems reasonable to implement adjustments at this level.

3 Working hours

Different exposure criteria were used in different studies, but in general studies considered working longer than around 40 hours, compared to 40 or less. Most of the data for these studies were collected after delivery.

One systematic review and nine original papers were identified that examined the relationship between working hours and the outcomes of interest.

Eight papers addressed preterm delivery, spontaneous miscarriage or perinatal mortality^{42,81,88,98,104,107,108,140} (five 2++ and three 2+) (Evidence tables 2 and 3). Five showed an increased risk of preterm birth with long working hours, with risk estimates of 1.24–1.7. Two showed an increased risk of spontaneous miscarriage, and one showed no effect with working >40 hours a week in previous pregnancies. A high quality systematic review²⁷ (1++) identified a further 11 papers (in addition to those in Evidence tables 2 and 3). Of these, eight were negative with relative risks close to unity, and five showed an increased risk of preterm delivery with relative risks of up to 1.87. However, the authors point out that the findings were statistically significant in only two of the four studies, where relative risks exceeded 1.5, and at least two studies had methodological limitations. Moreover, in the four largest studies (>2000 pregnancies analysed), relative risks were between 0.59 and 1.34. They carried out a meta-analysis, and found a pooled relative risk of 1.31 (95% CI 1.16 to 1.47, $Q=4.33$, $p=0.74$).

Two studies assessed low birth weight, IUGR or SGA in relation to long working hours^{108,161} (2+ and 2++) (Evidence tables 4 and 6). Both showed a positive relationship, with risk estimates 1.34 and 2.4. A high quality systematic review²⁷ (1++) identified a further six papers (in addition to those in Evidence tables 4 and 6), five of which were of high quality. The largest study found an increased risk (odds ratio 1.7), but the other five were all negative with relative risks close to unity.

One paper that examined pre-eclampsia was identified⁷³ (2+) (Evidence table 5). This study found a two-fold increase in the risk of pre-eclampsia (the study compared female residents with the spouses of male residents). A high quality systematic review²⁷ (1++) identified a further two papers (in addition to the paper in Evidence table 5). Both were of high quality and showed no relation between working hours and pre-eclampsia.

Conclusion

There was a large body of evidence for preterm delivery, spontaneous miscarriage/perinatal mortality. There was reasonable consistency, with 9/19 studies showing no effect and 10/19 showing risk estimates of no greater than 2.0. A high quality meta-analysis gave a pooled relative risk of 1.3. Overall the evidence suggested a low risk, and made more than a moderate risk unlikely. The evidence for low birth weight/IUGR/SGA was reasonable in volume and consistency, tending to suggest no effect or no more than a moderate effect. There was very limited evidence for pre-eclampsia.

RECOMMENDATIONS

Recommendation	Risk	Grade	Evidence
Pregnant employees should be informed about the generally consistent evidence suggesting that long working hours carry no more than small to moderate risk of preterm birth, and low birth weight/SGA, but there is limited inconsistent evidence for pre-eclampsia	No greater than a low risk of preterm birth, preterm labour or spontaneous miscarriage	A	42, 81, 88, 98, 104, 107, 108, 140, review ref 27 includes a further 11 papers
Employers should reduce long working hours for pregnant workers, particularly in late pregnancy. Working hours should be limited as far as possible to a 'standard' working week of approximate 40 hours per week. However, if a pregnant worker who has been informed of the possible risks wishes to continue then there are insufficient grounds upon which to impose restrictions against her will.	No more than a moderate risk of low birth weight		108,161, review ref 27 includes a further 6 papers
	Insufficient evidence for pre-eclampsia		81, review ref 27 includes a further 2 papers

Most studies about long working hours used >36 or >40 hours as a definition, therefore it seems reasonable to use 40 hours as the threshold for applying adjustments to work. It is unclear from current evidence at which stage of pregnancy this is advisable.

4 Shift work

This review identified nine studies and one systematic review that examined shift work or night work in relation to the outcomes of interest.

Seven studies addressed preterm birth, spontaneous miscarriage or perinatal mortality^{15,25,49,104,107,169,171} (six 2+ and one 2++) (Evidence tables 2 and 3). Of three that examined preterm birth, two found an increased risk with odds ratios 5.6 and 2.0. One found a risk of 1.45, but this was not statistically significant. Among four that looked at spontaneous miscarriage, one did not find a significant association; but three showed an increased risk, including one with high risk estimates of 2.7 for two-shift schedules and up to 6.89 for night work. A high quality systematic review²⁷ (1++) identified a further nine papers (excluding those in Evidence tables 2 and 3), all but one of which were high quality. In five, the risk estimates were close to one. Four showed an increase risk with risk estimates ≥ 1.5 . However, the authors pointed out that the positive studies tended to be small, and the four largest studies found little association between shift work and preterm delivery. The pooled estimate of risk across 13 studies was 1.2 (95% CI 1.01 to 1.42, $Q=31.3$, $p=0.002$).

Four studies were found that explored low birth weight^{14,17,15,169} (2+) (Evidence table 4). All four showed an increased risk of low birth weight, including three for rotating shifts (risk

estimate stated by one study was 2.5) and two for night or evening shifts (odds ratio in one study 5.6). A high quality systematic review²⁷ (1++) identified a further five papers (in addition to those in Evidence tables 4 and 6), including four of high quality. One of these showed an elevated risk with relative risk 1.5, but the rest showed risk estimates close to unity. In particular, a very large study of >35000 pregnancies showed no relationship between shift work and SGA. Moreover, the authors carried out a meta-analysis, which gave a pooled estimate of risk from six studies as 1.07 (95%CI 0.96 to 1.19, Q=3.30, p=0.51).

One study was identified that looked at pre-eclampsia¹⁶⁵ (2+) (Evidence table 5), finding a relative risk of 1.3 (although this was not statistically significant). A high quality systematic review²⁷ (1++) identified one further paper (in addition to the one in Evidence table 5). This showed no significant association (relative risk 0.9).

Conclusions

There was a reasonable body of evidence for shift work and preterm birth, but this was conflicting. Nine of the 16 papers identified (including those in a systematic review) showed an increased risk, including some with high relative risks of >2.0. However, seven studies had risk estimates close to one, and a high quality meta-analysis (that included four of the positive studies) showed a pooled estimate of 1.2. Similarly for low birth weight there was a reasonable volume of evidence but conflicting findings. About half of the studies gave positive and half negative results, although a meta-analysis supported no effect. There was insufficient evidence for pre-eclampsia (two papers).

There is insufficient evidence of a risk to pregnant women to make recommendations to restrict shift work, including rotating shifts or night/evening work.

5 Fatigue score

This score was devised by Mamelle⁹⁹ and is based upon five sources of occupational fatigue (posture, work on industrial machine, physical exertion, mental stress, and the working environment); Mamelle *et al* constructed an index based on these fatigue sources whereby a high score relates to increasing fatigue. In two studies a fatigue score ≥ 3 was related to a risk of preterm birth^{88,98} (2++). A dose-response relationship was found in the first study.⁹⁸ Other studies^{116,94} have found an association with each component of the fatigue score and preterm rupture of membranes, and adjusted odds ratios were significant for posture and physical exertion.

Overall, based on just three studies, there is insufficient evidence to make recommendations about the use of the fatigue score to manage pregnant women at work. In addition it is difficult to give advice on how a fatigue score would translate into work place restrictions.

4 Future research

The literature review on which these guidelines are based identified a substantial body of research into the workplace risk factors for adverse outcomes of pregnancy. However, the GDG has been unable to make recommendations for practice in some areas due to a relative paucity of evidence that is both of high quality and consistent. The main priorities for new research are specific exposures and pregnancy outcomes for which the current evidence is either small in volume (small number of papers) or where there is currently an important degree of inconsistency of results or serious methodological limitations.

The impact of working shift patterns and night work, particularly on measures of impaired fetal growth and pre-eclampsia, is an important priority area. At present the balance of evidence points to no effect or at most a low risk, but a small number of studies reported high risks (>2). As shift patterns are very common in some female-dominated jobs (eg nursing) more research would be helpful in guiding practice in this area.

The number of studies that addressed pre-eclampsia and pregnancy-induced hypertension were generally fewer than for the other outcomes, and this is another important priority area for further research. A systematic review²⁷ highlighted a problem with variation in the definition of pre-eclampsia (degree of proteinuria or level of blood pressure required for diagnosis) that would need to be addressed in future studies in this area.

The evidence base for preterm delivery and spontaneous miscarriage was more extensive than the other outcomes of interest, and (with the possible exception of heavy physical work) it was fairly consistent. However, a degree of inconsistency was seen for heavy physical work, with a moderate number of studies finding risk estimates >1.5. Moreover, there was considerable heterogeneity in exposure assessment. Therefore some more research into the consistent categorisation of heavy work and the association with preterm labour would be useful.

Finally, a common problem with the existing literature in this field is the reliance on retrospective assessment of exposure at the end of the pregnancy. In view of the important potential for bias when the outcome of pregnancy is known, there is a need for more prospective studies of the effect of work and working conditions on pregnancy outcome.

Recommendations for research

- (1) Prospective studies of the effect of shift work and night work on birth weight.
- (2) Prospective studies of the effect of workplace exposures on pre-eclampsia, with agreement of common definitions for outcome assessment.
- (3) Prospective studies of lifting and heavy physical activity, using consistent methods of exposure assessment and examining all adverse outcomes of pregnancy.

5 Audit criteria

Audit question	Standard
Is a risk assessment being undertaken by the employer upon notification of pregnancy?	100%
Is the pregnant employee given appropriate information about health risks in the workplace, including physical work and working hours?	100%
Is restriction of duties allowed if the pregnant worker requests it? And, if yes, are pay and conditions preserved?	100%

Evidence table 1 Summary of papers included as evidence

First author	Location	Study period	Study design	Exposure	Method of exposure assessment	Outcome(s)
Ahlborg (1990) ⁹	Orebro, Sweden	1980–3	Prospective cohort	Lifting	Self-administered questionnaire during pregnancy with follow-up after delivery	Preterm birth Low birth weight Spontaneous miscarriage
Armstrong (1989) ¹⁴	Montreal, Canada	1982–4	Retrospective cohort	Lifting Shift work		Low birth weight
Ali (2002) ¹¹	Peninsular Malaysia	2002	Case-control	Prolonged standing	Questionnaire	Perinatal mortality
Axelsson (1989) ¹⁷	Sweden	1980–4	Cross-sectional	Shift work	Mailed questionnaire	Low birth weight
Axelsson (1996) ¹⁵	Sweden	1989	Retrospective cohort	Shift work in first trimester	Mailed questionnaire	Spontaneous miscarriage
Bodin (1999) ²⁵	Sweden	1980–7	Retrospective cohort	Shift work	Mailed questionnaire	Low birth weight Preterm birth
Bonzini (2007) ²⁷	Worldwide	1966–2005	Systematic review and meta-analysis	Working hours Shift work Lifting Standing Physical activity	Systematic review of literature	Preterm delivery Small-for-gestational-age Pre-eclampsia/hypertension
Eskenazi (1994) ⁴²	California, USA	1986–7	Case-control	Physical exertion Heavy lifting Working hours	Questionnaire after delivery	Spontaneous miscarriage
Fenster (1997) ⁴⁶	California, USA	1990–1	Case-control and part of prospective cohort	Standing	Interview using computer-assisted telephone interview system	Spontaneous miscarriage
Fortier (1995) ⁴⁹	Quebec city, Canada	1989	Retrospective cohort	Shift work Working hours Standing Lifting Physical activity	Telephone interview after delivery	Preterm delivery Small-for-gestational-age
Ha (2002) ⁵⁸	Beijing, China	1996–8	Prospective cohort	Physical activity Standing	Interview using structured questionnaire	Low birth weight
Henriksen (1995) ⁶⁴	Netherlands	1989–91	Prospective cohort	Standing Walking	Self-administered questionnaire at two stages of pregnancy	Preterm birth
Henriksen (1995) ⁶⁴	Denmark	1989–91	Prospective cohort	Standing Walking	Self-administered questionnaire at 16 weeks and 30 weeks gestation	Low birth weight

continued

Physical and shift work in pregnancy: occupational aspects of management

First author	Location	Study period	Study design	Exposure	Method of exposure assessment	Outcome(s)
Infante-Rivard (1993) ⁷⁴	Canada	1987–9	Case-control	Shift work	Face-to face interview post delivery	Spontaneous miscarriage
Kaerlev (2004) ⁷⁷	Denmark	1995–9	Retrospective cohort	Working hours Shift work Lifting Standing Walking	Self-administered questionnaire	Sick leave
Klebanoff (1990) ⁸¹	USA	1985	Retrospective cohort and case-control	Working hours Shift work	Mailed questionnaire	Preterm labour Pre-eclampsia
Koemeester (1995) ⁸³	Netherlands	1989–90	Prospective cohort	Physical workload	Questionnaire and interview at various stages of pregnancy	Gestational age
Lima (1999) ⁸⁶	Brazil	April–December 1992	Retrospective cohort	Heavy work	Interviews in hospital after delivery	Low birth weight
Luke (1995) ⁸⁸	USA	1991	Case-control	Working hours Occupational fatigue (defined by fatigue score)	Mailed structured questionnaire	Preterm birth
Mamelle (1984) ⁹⁸	France	1977–8	Retrospective cohort	Working hours Standing Occupational fatigue (defined by fatigue score)	Questionnaire based interview immediately after delivery	Preterm birth
McDonald (1986) ¹⁰⁴	Montreal, Canada	1982–4	Retrospective cohort	Lifting Standing Working hours Shift work	Questionnaire based interview soon after delivery	Spontaneous abortion
McDonald (1988) ¹⁰⁷	Montreal, Canada	1982–4	Retrospective cohort	Lifting Standing Physical activity Working hours	Questionnaire based interview soon after delivery	Spontaneous abortion
McDonald (1988) ¹⁰⁸	Montreal, Canada	1982–4	Retrospective cohort	Lifting Working hours Physical effort Standing Shift work	Questionnaire based interview soon after delivery	Preterm birth Low birth weight
Newman (2001) ¹¹⁶	USA		Prospective cohort	Occupational fatigue (defined by fatigue score)	Nurse-administered standard questionnaire at enrolment; follow-up questionnaire between 27 and 31 weeks	Preterm birth
Ritsmitchai (1997) ¹³³	Thailand	1993	Case-control	Standing Physical exertion	Interview using structured questionnaire	Preterm birth

continued

Evidence table 1 Summary of papers included as evidence

First author	Location	Study period	Study design	Exposure	Method of exposure assessment	Outcome(s)
Saurel-Cubizolles (1987) ¹³⁸	France	1981	Cross-sectional study taken from a national survey	Physical effort Standing and carrying a heavy load	Interview using a standardised questionnaire within six days of delivery	Preterm birth Low birth weight
Saurel-Cubizolles (1991) ¹⁴¹	France	1987–8	Retrospective survey	Standing Lifting Working hours	Structured questionnaire after delivery in hospital	Preterm birth
Saurel-Cubizolles (2004) ¹⁴⁰	17 European countries	1994–7	Case-control	Standing Working hours	Interview using questionnaire after delivery	Preterm birth
Spinillo (1996) ¹⁵²	Italy	1989–94	Case-control	Physical effort	Structured interview at birth	Fetal growth restriction
Strand (1997) ¹⁵⁵	Norway	1989	Cross-sectional	Physical effort Lifting Working hours Work schedules	Questionnaire after delivery in hospital	Sick certification
Teitelman (1990) ¹⁵⁹	USA	1980–2	Prospective cohort	Standing	Interview using a standard questionnaire in first trimester (and further 20%) by 20 weeks	IUGR Preterm birth
Tuntiseranee (1998) ¹⁶¹	Thailand	1994–5	Prospective cohort	Working hours Lifting	Interviews at weeks 17 and 32	Small-for-gestational-age Low birth weight Preterm delivery
Wergeland (1997) ¹⁶⁵	Norway	1989	Retrospective cohort	Lifting Work above shoulder level Shift work	Questionnaire completed after delivery	Pre-eclampsia
Xu (1994) ¹⁶⁹	China	1992	Retrospective cohort	Shift work	Standardised questionnaire	Preterm birth Low birth weight
Zhang (1996) ¹⁷⁰	Connecticut, USA	1988–91	Prospective cohort	Manual handling – reaching above shoulders Standing	Home interview between 5 and 16 weeks gestation	Spontaneous abortion
Zhu (2004) ¹⁷²	Denmark	1998–2001	Prospective cohort	Shift work	Computer assisted telephone interviews twice during pregnancy and twice after delivery. Face-to-face interview post delivery	Duration of pregnancy
Zhu (2004) ¹⁷¹	Denmark	1998–2001	Prospective cohort	Shift work	Telephone interviews between 11 and 25 weeks, again around 30 weeks, and post delivery	Spontaneous miscarriage

OR = odds ratio; RR = relative risk

Evidence table 2 Preterm birth/threatened preterm labour

Author	Study population	Main results	Effect measure	Exposure comparison and timing in pregnancy	Confounders	Sign grade
Lifting						
<i>Systematic review</i>						
Bonzini (2007) ²⁷	Association between lifting and preterm birth studied in 12 papers	Substantial difference in definitions of exposure. RR \leq 1.35 in 11 of 12 studies, RR \geq 1.5 in none of the 12 studies	See previous column	Various, mainly first trimester	Judged low risk of confounding in 10 of 12 studies	1++
<i>Prospective cohort</i>						
Ahlborg (1990) ⁹	All pregnant women in antenatal care centres in Örebro county in the study period (1980–3) who worked during pregnancy, n=3906 (based on the number of subjects in the final analysis)	No significant risk associated with heavy lifting and preterm birth.	Adjusted OR 1.29 (0.69–2.4) (Table 4)	12 kg or more for more than 50 times per week (heaviest exposure) vs none first trimester. Exposure data were collected during early pregnancy, (from the time of LMP until first visit to antenatal care.	Age, parity, sex of infant, educational level, smoking and frequency of alcohol use	2+
<i>Retrospective cohort</i>						
McDonald (1988) ¹⁰⁸	Women attending 11 obstetrical units in Montreal from 1982–4, n=51,885, n=56,067 interviewed for the study. Results based upon n=2,2761	Significant association between lifting heavy weights and preterm birth	O/E 1.26 p<0.01	Lifting heavy weights \geq 15 vs <15 times per day. Timing not noted	Age, gravidity, previous spontaneous abortion, ethnic group, height, educational level, cigarette smoking and alcohol consumption	2++

continued

Evidence table 2 Preterm birth/threatened preterm labour

Author	Study population	Main results	Effect measure	Exposure comparison and timing in pregnancy	Confounders	Sign grade
Manual handling						
<i>Prospective cohort</i>						
Tuntiseranee (1998) ¹⁶¹	Pregnant women attending routine antenatal care at the dept of Obstetrics in 2 Thai hospitals from 1994–1995, n=1821	No significant risk of preterm delivery associated with lifting	RR 0.9 (0.4–2.1)	Lifting >12 kg 1–10 times per day vs none. Timing 15–28 weeks	Parity, maternal height, maternal age, obstetrical complications, baby sex, number of antenatal visits, maternal weight at delivery	2+
Physical activity						
<i>Prospective cohort</i>						
Koemeester (1995) ⁸³	Nurses working in 12 hospitals between 1989–90 who worked for at least the first 12 weeks of pregnancy, n=116 (number in the study)	There was a significant association between gestational age at delivery and duration of high physical workload. After stepwise multiple regression analysis on nurses without extra risks in pregnancy, the daily duration of tasks with a high physical workload, was significantly correlated with shorter gestational age at delivery	p=0.01 p=0.004	Daily duration of tasks with high physical workload in 3 categories (<2 hours, 2–4 hours, >4 hours). Timing not noted	Smoking, drinking, parity, age, sporting activities, non-physical occupational factors, work hours per week, frequency of stooping per day, frequency of lifting per day, frequency of standing per day duration of tasks with physical workload (stepwise multiple regression analysis)	2+
<i>Retrospective survey</i>						
Saurel-Cubizolles (1991) ¹⁴¹	Random sample of births in four public maternity units in France between April 1987 and May 1988, n=1949 interviewed final sample from which results obtained n=875	Significant difference in preterm births between different occupations. There was a difference in the physical working conditions of different occupations. However there was no significant difference in preterm births according to working conditions. Of note: women with strenuous working conditions reported more often that they had changed their working conditions. These women also stopped work before 28 weeks	RR 1.31 (0.64–2.58)	Lifting heavy loads often/always vs none/sometimes. Timing not noted	Age, nationality, education level, gravidity, previous preterm birth, maternity centre	2+

continued

Physical and shift work in pregnancy: occupational aspects of management

Author	Study population	Main results	Effect measure	Exposure comparison and timing in pregnancy	Confounders	Sign grade
Physical activity – continued						
<i>Case-control</i>						
Ritsmitchai (1997) ¹³³	Mothers from Hat Yak regional hospital in 1993 Cases: spontaneously born singleton preterm infants delivered between 28 and 37 weeks, n=223 Controls: the first mothers of singleton infants delivered at over 37 weeks after each case – matched, n=223	Significant risk of preterm birth if physical exertion throughout pregnancy	Adjusted OR 2.91 95% CI 1.29–6.58	All pregnancy	Pregnancy complications, previous preterm birth, physical exertion, physical exercise. Did not adjust for coffee, alcohol, smoking, because small numbers exposed	2+
<i>Cross-sectional from a national survey</i>						
Saurel-Cubizolles (1987) ¹³⁸	Randomly sampled women who delivered in French maternity units in 1981 n=5508 interviewed study based upon n=2387	Higher rate of preterm birth among women who had to make considerable physical effort, and women who cumulated >2 strenuous working conditions. Not significant after adjustment for occupation. (note – is appropriately in table 4 as well) Significant increase in preterm delivery if undertaking 1 or 2 of the strenuous working conditions	p=0.05	Heavy load carrying yes vs no. Considerable physical effort yes vs no Timing: first trimester. The proportion of women who did not work in the third trimester was higher when the work was strenuous. Also more likely to change jobs	Parity, smoking, preterm delivery	2+
Standing						
<i>Systematic review</i>						
Bonzini (2007) ²⁷	20 papers considered standing and preterm birth	‘High’ exposure was defined as >3 hours in 12 studies. RR>1.5 in seven studies, less clearly positive in the rest. Meta-analysis RR=1.28	Pooled estimate for 12 studies RR 1.28 95% CI 1.11–1.47	See references in review	See references in review	1++

continued

Evidence table 2 Preterm birth/threatened preterm labour

Author	Study population	Main results	Effect measure	Exposure comparison and timing in pregnancy	Confounders	Sign grade
Standing – continued						
<i>Prospective cohort</i>						
Teitelman (1990) ¹⁵⁹	Eligible women who sought care from various health care providers in New Haven, expected to deliver at Yale-New Haven hospital and were pregnant, between 1980 and 1982; n=6219 introduced to the study; n=4186 completed valid interviews; final analysis based upon n=1,206	Significant risk of preterm delivery in those standing >3 hours/day, compared with sedentary and active jobs	Adjusted OR 2.72 95% CI 1.24–5.95	Not noted	Parity, smoking, education, caffeine use, illicit drug use, gestational age at interview and marital status	2+
<i>Retrospective cohort</i>						
Mamelle (1984) ⁹⁸	Women giving birth in 2 hospitals between 1977–8, analysis based upon n=1928 working women	Significantly increased risk if standing >3 hours	RR 1.6 95% CI 1.1–2.3	Not noted	Low socio-economic level, unmarried mother, low maternal education level, low maternal age, primiparity, previous prematurity, obstetric pathology	2++
<i>Case-control</i>						
Ritsmitchai (1997) ¹³³	Mothers from Hat Yai regional hospital in 1993 Cases: spontaneously born singleton preterm infants delivered between 28 and 37 weeks n=223 Controls: the first mothers of singleton infants delivered at over 37 weeks after each case – matched, n=223	Cases were more likely than controls to have been exposed to prolonged standing in all three trimesters. Significant risk of preterm birth if standing >3 hours/day throughout pregnancy	Adjusted OR 4.10 95% CI 1.29–13.10 p=0.001	Not noted	Pregnancy complications, previous preterm birth, physical exertion, physical exercise. Did not adjust for coffee, alcohol, smoking, because small numbers exposed	2+

continued

Author	Study population	Main results	Effect measure	Exposure comparison and timing in pregnancy	Confounders	Sign grade
Standing – continued						
<i>Case-control – continued</i>						
Saurel-Cubizolles (2004) ¹⁴⁰	Cases: Consecutive singleton pre-term births in 17 countries between 1994–1997, working after 3 rd month of pregnancy n=2369 Controls randomly selected (not matched) n=4098	Significantly higher risk of preterm birth if standing >6 hours/day	Adjusted OR 1.26 95% CI 1.1–1.5 In A2 countries: Adjusted OR 1.38 95% CI 1.1–1.7 In B countries: Adjusted OR 1.55 95% CI 1.1–2.3	Not noted	Maternal age, women’s educational level, marital status, obstetric history	2++
Standing and walking						
<i>Prospective cohort</i>						
Henriksen (1995) ⁶⁵	Women attending for antenatal care at Aarhus university hospital from 1989–91, analysis based upon n=4259	Significantly greater risk of preterm delivery if walking and standing for >5 hours/day (does not distinguish between the two), compared with 2 hours or less. Majority still working after 30 weeks, but a significant number were either off sick or no longer employed	Adjusted RR 3.3 95% CI 1.4–8.0	More women who reported jobs involving long hours of standing and walking at 16 weeks, were no longer employed by the 30 th week. Those who left jobs involving standing or walking prior to 30 weeks had the same risk of preterm deliveries as women with the same exposure in the 16 th week who kept working through the third trimester	Parity, maternal height, smoking, leisure time activities, partner’s social class	2+
Working hours						
<i>Systematic review</i>						
Bonzini (2007) ²⁷	16 papers considering the relationship between working hours and preterm delivery	Moderate associations were reported. RR >1.5 in four studies, but some limitations. Lower RR in the larger studies	Eight studies pooled estimate: RR 1.31 95% CI 1.16–1.47	See references in review	See references in review	1++

continued

Evidence table 2 Preterm birth/threatened preterm labour

Author	Study population	Main results	Effect measure	Exposure comparison and timing in pregnancy	Confounders	Sign grade
Working hours – continued						
<i>Retrospective cohort</i>						
Klebanoff (1990) ⁸¹	Women who graduated from medical school in 1985 and a random sample of male physician spouses Final study population: 989 residents, 1239 spouses	Residents worked >64 hours/week compared with <37 hours/week in spouses. Twice as many women residents going in to threatened preterm labour requiring bed rest or hospitalisation	p<0.001	Not noted	Parity, age at delivery, height, pre-pregnancy weight and race or ethnic group	2+
Mamelle (1984) ⁹⁸	Women giving birth in 2 hospitals between 1977–8, analysis based upon n=1928 working women	Significantly increased risk of prematurity if working >41 hours/week. Note, increasing % of prematurity with increasing hours worked	RR 1.7 95% CI 1.1–2.5	Not noted	Low socio-economic level, unmarried mother, low maternal education level, low maternal age, primiparity, previous prematurity, obstetric pathology	2++
McDonald (1988) ¹⁰⁸	Women attending 11 obstetrical units in Montreal from 1982–4, n=56067 interviewed for the study. Results based upon n=22761	Significant association between working >46 hours/week and preterm birth	O/E 1.24 p=0.03	Not noted	Age, gravidity, previous spontaneous abortion, ethnic group, height, educational level, cigarette smoking and alcohol consumption	2++
<i>Case-control</i>						
Luke (1995) ⁸⁸	Cases: Nurses whose infants were delivered preterm (<37 weeks) in 1991 with current membership of The Association of Women's Health, Obstetrics and Neonatal Nurses (AWHONN), n=210 Controls: nurses whose infants were delivered at term (>37 weeks), n=1260	Significantly increased risk of preterm birth if working >36 hours/week, compared with ≤36 hours/week	Adjusted OR 1.6 95% CI 1.1–2.2 p=0.006	Not noted	Young maternal age, non-white race, unmarried status, low educational level, pregnancy complications	2++

continued

Physical and shift work in pregnancy: occupational aspects of management

Author	Study population	Main results	Effect measure	Exposure comparison and timing in pregnancy	Confounders	Sign grade
Working hours – continued						
<i>Case-control</i>						
Saurel-Cubizolles (2004) ¹⁴⁰	Cases: Consecutive singleton preterm births in 17 countries between 1994–7, working after 3 rd month of pregnancy n=2369 Controls randomly selected (not matched) n=4098	Significantly higher risk of preterm birth if working >43 hours/week	Adjusted OR 1.33 95% CI 1.1–1.6	Not noted	Maternal age, women’s educational level, marital status, obstetric history	2++
Shift work						
<i>Systematic review</i>						
Bonzini (2007) ²⁷	14 studies considered the association between preterm delivery and shift work	In 7 studies the RR was close to unity; in 6 studies the RR was >1.5. Two studies found risks >2.0. Meta-analysis RR=1.2	Pooled estimates of risk in 13 studies: RR 1.20 95% CI 1.01–1.42	See references in review	See references in review	1++
<i>Retrospective Cohort</i>						
Bodin (1999) ²⁵	Female members of the Swedish Midwives Association in 1989; pregnancies between 1980–1987, n=1244 number included in analysis	Significant association between night work and preterm birth	Adjusted OR 5.6 95% CI 1.9–16.4	Not noted	Maternal age, parity, employment, work schedule, N2O use	2+
Fortier (1995) ⁴⁹	Women in 6 community health departments in Quebec city who gave birth to a live singleton in 1989 Analysis based upon n=4390	For women working regular evening or night work compared with women always working days, the adjusted OR of preterm birth was 1.45x higher, however, not statistically significant	See previous column	Not noted	Age, parity, history of low birth weight or preterm delivery, education, family income, pre-pregnancy weight, height, alcohol and caffeine intake, active and passive smoking, and energy expenditure in non-occupational activities	2++

continued

Evidence table 2 Preterm birth/threatened preterm labour

Author	Study population	Main results	Effect measure	Exposure comparison and timing in pregnancy	Confounders	Sign grade
Shift work – continued						
<i>Retrospective Cohort</i>						
Xu (1994) ¹⁶⁹	Married women working in textile mills in China. Analysis based upon n=887 live births. Married, educated, non-smokers, non-alcohol drinkers	Significant increase in prevalence of preterm birth of all live births associated with rotating shifts.	Adjusted OR 2.0 95% CI 1.1–3.4 p=0.02	Not noted	Mill location, maternal age at pregnancy, job title, time and duration of leave from the job since pregnancy, order of live births, exposure to dust/gases/fumes, physical activity and position while working, stress at work and use of coal stove for heating at home	2+
Occupational fatigue						
<i>Prospective cohort</i>						
Newman (2001) ¹¹⁶	Cohort of 2929 singleton pregnancies part of the multicentre preterm prediction study. Analysis based upon this number	Each of the 5 sources of occupational fatigue were associated with a significant increased risk of preterm premature rupture of membranes in nulliparous women. Significant linear trend (p=0.002) between risk of preterm premature rupture of membranes in nulliparous women and increasing occupational fatigue index. Significant association between hours worked per week and preterm premature rupture of membranes (p=0.01). With multivariate logistic regression, the adjusted odds ratio showed significant association of each individual source of occupational fatigue and overall occupational fatigue score for preterm premature rupture of membranes in nulliparous women	Posture adjusted OR 1.69 95% CI 1.20–2.38 Physical exertion OR 1.72 95% CI 1.16–2.23	Not noted	Contractions, cervical length, BMI, race, living with another adult, education, private insurance, smoking during pregnancy, alcohol use during pregnancy, drug use during pregnancy, vaginal bleeding	2+

continued

Physical and shift work in pregnancy: occupational aspects of management

Author	Study population	Main results	Effect measure	Exposure comparison and timing in pregnancy	Confounders	Sign grade
Occupational fatigue – <i>continued</i>						
<i>Retrospective cohort</i>						
Mamelle (1984) ⁹⁸	Women giving birth in two hospitals between 1977–8, analysis based upon n=1928 working women	Risk of prematurity increases when a woman accumulates two or more sources of occupational fatigue	High fatigue score with no medical factor present: RR 2.1 95% CI 1.3–3.5 With medical factor present RR 2.8 95% CI 1.8–4.9	Not noted	Low socio-economic level, unmarried mother, low maternal education level, low maternal age, primiparity, previous prematurity, obstetric pathology	
<i>Case-control</i>						
Luke (1995) ⁸⁸	Cases: nurses whose infants were delivered preterm (<37 weeks) in 1991 with current membership of AWHONN, n=210 Controls: nurses whose infants were delivered at term (>37 weeks), n=1260	Significantly increased risk of preterm birth if occupational fatigue score ≥ 3 . Dose response relationship between increasing fatigue score and increasing hours of work. Note the higher the occupational fatigue score, the more tired the nurse at the end of the shift. Highest fatigue scores in acute clinical areas	Adjusted OR 1.4 95% CI 1.1 –1.9 p=0.023 Cumulative effect: Adjusted OR 2.2	Not noted	Young maternal age, non-white race, unmarried status, low educational level, pregnancy complications	2++

OR = odds ratio; RR = relative risk

Evidence table 3 Spontaneous miscarriage/perinatal mortality

Author	Study population	Main results	Effect measure	Timing in pregnancy	Confounders	Sign grade
Lifting						
<i>Prospective cohort</i>						
Ahlborg (1990) ⁹	All pregnant women in antenatal care centres in Örebro county in the study period (1980–3) who worked during pregnancy, n=3906 the number used in the final analysis	No significant risk associated with heavy lifting and fetal death. The heaviest work categorised as >12 kg >50 times a week	Adjusted OR 1.06 95% CI 0.62–1.81	Lifting in first trimester	Age, parity, sex of infant, educational level, smoking and frequency of alcohol use	2+
<i>Retrospective cohort</i>						
McDonald (1986) ¹⁰⁴	Women attending 11 obstetrical units in Montreal from 1982–4, n=56,012 interviewed n=4127 + 10,910 met the criteria for spontaneous abortion	Ratios of O/E miscarriages significantly raised for early, mid and late miscarriages if lifting heavy weights >15 times daily. Significant increase in spontaneous miscarriage if lifting heavy weights in various occupational sectors (Table 2 of paper). A significant trend of increasing risk of spontaneous miscarriage with increasing amount of heavy lifting, for current pregnancies (p<0.01) and previous pregnancies (p<0.01)	Heavy lifting >15 times: RR 2.00 95% CI 1.5–2.5	Not noted	Maternal age, parity, history of previous abortion, smoking habit, highest education level reached	2+
McDonald (1988) ¹⁰⁷	Women attending 11 obstetrical units in Montreal from 1982–4; n=56067 interviewed n=22,613 previous pregnancies meeting criteria, therefore number used in analysis	Significantly increased risk of spontaneous miscarriage before 28 weeks associated with lifting heavy weights >15 times a day. For heavy lifting the O/E ratio increased with increasing grade of the factor	RR 2.0 90% CI 1.5–2.5 p<0.01 (p<0.01)	Not noted	Age, gravidity, previous spontaneous abortion, ethnic group, height, educational level, cigarette smoking and alcohol consumption	2+

continued

Physical and shift work in pregnancy: occupational aspects of management

Author	Study population	Main results	Effect measure	Timing in pregnancy	Confounders	Sign grade
Lifting – continued						
<i>Case-control</i>						
Eskenazi (1994) ⁴²	Cases: pregnant women in Santa Clara County who had spontaneous miscarriage before 20 weeks gestation, from 1986–7, n=434 (working) Controls: two per case randomly selected from a matched group, n=910 (working)	No evidence of association with spontaneous miscarriage and frequently lifting >15 lbs	Adjusted OR 1.1 95% CI 0.6–2.0	Not noted	Race, maternal age history of spontaneous miscarriage, cigarette smoking, alcohol, caffeine consumption, tap water consumption, marital status, parity, level of education and nausea	2+
Manual handling						
<i>Prospective cohort</i>						
Zhang (1996) ¹⁷⁰	Women receiving prenatal care in organizations in Connecticut (Oct 1988–Dec 1991) who had a singleton live birth or a spontaneous abortion, n=2849 number used in analysis	Significant risk if carrying loads >9 kg once or more a day. Reaching over the shoulders once or more a day	Adjusted RR 1.71 95% CI 1.25–2.32 Adjusted OR 1.75 95% CI 1.13–2.71 Adjusted RR 1.35 95% CI 1.02–1.78	First trimester	Those considered – not all used at every stage: maternal age, marital status, mother’s height, smoking, illicit drug use, chronic medical problems, use of birth control, number of pregnancies, induced abortion, spontaneous miscarriage, education, race, stopped smoking before pregnancy, infertility, stillbirth, ectopic pregnancy	2+
Physical activity						
<i>Retrospective cohort</i>						
McDonald (1988) ¹⁰⁷	Women attending 11 obstetrical units in Montreal from 1982–4; n=56,067 interviewed; n=22,613 previous pregnancies meeting criteria, therefore number used in analysis	Significantly increased risk of spontaneous miscarriage before 28 weeks associated with physical effort. For physical effort O/E ratio increased with increasing grade of the factor	RR 1.87 90% CI 1.4–2.3 p<0.01 (p<0.01)	Not noted	Age, gravidity, previous spontaneous miscarriage, ethnic group, height, educational level, cigarette smoking and alcohol consumption	2++

continued

Evidence table 3 Spontaneous miscarriage/perinatal mortality

Author	Study population	Main results	Effect measure	Timing in pregnancy	Confounders	Sign grade
Standing						
<i>Retrospective cohort</i>						
McDonald (1986) ¹⁰⁴	Women attending 11 obstetrical units in Montreal from 1982–4, n=56,012 interviewed n=4127 + 10,910 met the criteria for spontaneous abortion	Ratios of O/E miscarriages significantly raised for early, mid and late miscarriages if standing >8 hours daily; significantly greater O/E ratio for stillbirths if standing >8 hours daily	Standing >8 hours daily: RR 1.32 95% CI 1.1–3.5	Not noted	Maternal age, parity, history of previous miscarriage, smoking habit, highest education level reached	2+
McDonald (1988) ¹⁰⁷	Women attending 11 obstetrical units in Montreal from 1982–4; n=56067 interviewed n=22,613 previous pregnancies meeting criteria, therefore number used in analysis	Significantly increased risk of spontaneous miscarriage before 28 weeks associated with standing >8 hours/day: (See Table 5 for breakdown by early, mid and late miscarriage.) Significant risk of stillbirth (>28 weeks) with standing >8 hours daily	RR 1.32 90% CI 1.1–3.5 p<0.01	Not noted	Age, gravidity, previous spontaneous miscarriage, ethnic group, height, educational level, cigarette smoking and alcohol consumption	2++
<i>Case-control</i>						
Eskenazi (1994) ⁴²	Cases: pregnant women in Santa Clara County who had spontaneous miscarriage before 20 weeks gestation, from 1986–7, n=434 (working) Controls: two per case randomly selected from a matched group, n=910 (working)	There is a significant association between standing >8 hours/day at work and spontaneous miscarriage. In second trimester: risk is much greater if there is a history of spontaneous miscarriage.	Adjusted OR 1.6 95% CI 1.1–2.3 Adjusted OR 2.1 95% CI 1.2–3.5	Not noted	Race, maternal age history of spontaneous miscarriage, cigarette smoking, alcohol, caffeine consumption, tap water consumption, marital status, parity, level of education and nausea	2+
Fenster (1997) ⁴⁶	Eligible members of prepaid healthcare plan in three areas of California, from 1990–1, n=5342 interviewed Working: n=4064 Numbers used in analysis	Significant increased risk of spontaneous miscarriage if standing for >7 hours in women with a history of >2 previous spontaneous miscarriages	Adjusted OR 4.32 95% CI 1.59–11.72	Not noted	Maternal age, gestational age at interview, pregnancy history, smoking, alcohol and caffeine consumption and marital status	2+

continued

Physical and shift work in pregnancy: occupational aspects of management

Author	Study population	Main results	Effect measure	Timing in pregnancy	Confounders	Sign grade
Standing – continued						
<i>Case-control</i>						
Ali (2002) ¹¹	Cases: working mothers experiencing perinatal death. n=78 Controls: working mothers with a live baby, not resulting in early neonatal death, n=78	Prolonged standing >3 hours is significantly associated with perinatal mortality	OR 3.3 95% CI 1.7–6.5 p=0.001 Adjusted OR 3.5 95% CI 1.4–8.3	Not noted	Father’s smoking status, birth weight of baby, baby’s maturity at birth, past history of preterm delivery, mother’s condition at work in relation to perinatal death. Maternal age, gravidity, antenatal care, previous obstetric history, medical complications, baby’s gender, shift work, total hours of work per week, workplace hazards	2+
Working hours						
<i>Case-control</i>						
Eskenazi (1994) ⁴²	Cases: pregnant women in Santa Clara County who had spontaneous abortion before 20 weeks gestation, from 1986–1987, n=434 (working) Controls: two per case randomly selected from a matched group, n=910 (working)	No evidence of association with spontaneous miscarriage and working >40 hours/week	Adjusted OR 1.2 95% CI 0.8–1.9	Not noted	Race, maternal age history of spontaneous miscarriage, cigarette smoking, alcohol, caffeine consumption, tap water consumption, marital status, parity, level of education and nausea	2+
<i>Retrospective cohort</i>						
McDonald (1986) ¹⁰⁴	Women attending 11 obstetrical units in Montreal from 1982–4, n=56,012 interviewed n=4127 + 10,910 met the criteria for spontaneous miscarriage	A significant trend of increasing risk of spontaneous miscarriage with working >40 hours/week in previous pregnancies	(p<0.01)	Not noted	Maternal age, parity, history of previous miscarriage, smoking habit, highest education level reached	2+

continued

Evidence table 3 Spontaneous miscarriage/perinatal mortality

Author	Study population	Main results	Effect measure	Timing in pregnancy	Confounders	Sign grade
Working hours – continued						
<i>Retrospective cohort</i>						
McDonald (1988) ¹⁰⁷	Women attending 11 obstetrical units in Montreal from 1982–1984; n=56067 interviewed; n=22,613 previous pregnancies meeting criteria, therefore number used in analysis	For long hours of work, the O/E ratio increased with increasing grade of the factor	(p<0.01)	Not noted	Age, gravidity, previous spontaneous miscarriage, ethnic group, height, educational level, cigarette smoking and alcohol consumption	2++
Shift work						
<i>Prospective cohort</i>						
Zhu (2004) ¹⁷¹	The Danish National Birth Cohort – all pregnant women who spoke Danish well enough; 1998–2001, initial n=54,954 enrolled n=42,687 with a job and work schedule	Increased risk of late fetal loss with fixed night work, but not significant	Adjusted hazard ratio 1.81 95% CI 0.88–3.72	Not noted	Maternal age, gravidity, history of spontaneous miscarriage, smoking, pre-pregnancy BMI, occupation, working posture, working hours per week, heavy lifting, perceived physically strenuous work, support from co-workers	2+
<i>Retrospective cohort</i>						
Axelsson (1996) ¹⁵	Female members of the Swedish Midwives Association in 1989; pregnancies during which the woman worked as a midwife, n=1717 number used in analysis	When looking at first pregnancies only (n=428), there is a significantly increased risk of spontaneous miscarriage for night-work, and two-shift schedules Significantly increased risk of late spontaneous abortion with night work	Adjusted OR 6.89 95% CI 1.43–33.3 Adjusted OR 2.70 95% CI 1.09–6.72 Adjusted OR 3.33 95% CI 1.13–9.87	Exposure in first trimester only	Calendar year, age, previous spontaneous miscarriage, infection, full-time work, working time, use of N2O, shortage of staff	2+
McDonald (1986) ¹⁰⁴	Women attending 11 obstetrical units in Montreal from 1982–4, n=56,012 interviewed n=4127 + 10,910 met the criteria for spontaneous abortion	Greater O/E ratio of early and mid abortions if changing shift pattern. Significant increase in spontaneous miscarriage if rotating shift work, in various occupational sectors (Table 2 of paper)	(Table 2 of paper)	Not noted	Maternal age, parity, history of previous abortion, smoking habit, highest education level reached	2+

continued

Physical and shift work in pregnancy: occupational aspects of management

Author	Study population	Main results	Effect measure	Timing in pregnancy	Confounders	Sign grade
Shift work – continued						
<i>Case-control</i>						
Infante-Rivard (1993) ⁷⁴	Women attending a particular hospital with a diagnosis of spontaneous miscarriage or fetal death from 1987–1989; matched controls Cases n=331 Controls n=993	Significantly increased risk of pregnancy loss if work fixed evening shift. Majority ≤17 weeks gestation. 26% >17 weeks	Adjusted OR 4.17 95% CI 2.19–7.92	Not noted	Age, level of schooling, presence of children at home, coffee drinking and uterine anatomical abnormality	2+

OR = odds ratio; RR = relative risk

Evidence table 4 Low birth weight

Author	Study population	Main results	Effect measure	Timing in pregnancy	Confounders	Sign grade
Lifting						
<i>Prospective cohort</i>						
Ahlborg (1990) ⁹	All pregnant women in antenatal care centres in Örebro county in the study period (1980–3) who worked during pregnancy, n=3906 the number used in the final analysis	Indication that heavy lifting might affect weight, but not conclusive, if woman continued to work at least until the 32 nd week of pregnancy. Lifting weights and frequencies were defined	Adjusted OR 0.65 95% CI 0.24–1.77 for heaviest lifting category	Lifting in first trimester	Age, parity, sex of infant, educational level, smoking and frequency of alcohol use	2+
Tuntiseranee (1998) ¹⁶¹	Pregnant women attending routine antenatal care at the dept of Obstetrics in 2 Thai hospitals from 1994–5, n=1821 number analysed in study n=2084 initially recruited	Significantly increased risk of low birth weight if lifting at chest level	Adjusted OR 3.5 95% CI 1.4–8.3	Reduced physical activity if at work at 32 weeks	Parity, maternal height, maternal age, obstetrical complications, baby sex, number of antenatal visits, maternal weight at delivery. Not necessarily applicable to UK context	2+
<i>Retrospective cohort</i>						
Armstrong (1989) ¹⁴	Women attending 11 obstetrical units in Montreal from 1982–4, n=22,404 number in analysis. Further analysis from a previous study	Lifting heavy weights more than 15 times a day was significantly associated with low birth weight before and after accounting for gestational age	Mean per cent predicted birth weight for age 99.1% 90% CI: 98.3–99.9 p<0.05	Not noted	In previous paper, and then adjusted for occupational sector	2+
McDonald (1988) ¹⁰⁸	Women attending 11 obstetrical units in Montreal from 1982–4, n=56,067 interviewed for the study. Results based upon n=22,761	Significant association between lifting heavy weights and low birth weights	O/E 1.25 p<0.01	Not noted	Age, gravidity, previous spontaneous miscarriage, ethnic group, height, educational level, cigarette smoking and alcohol consumption	2++

continued

Physical and shift work in pregnancy: occupational aspects of management

Author	Study population	Main results	Effect measure	Timing in pregnancy	Confounders	Sign grade
Physical activity						
<i>Retrospective cohort</i>						
Lima (1999) ⁸⁶	Women with low-income, with babies born April–December 1992 in NE Brazil; n=958 number in analysis	Significant reduction in birth weight if did heavy agricultural work for nine months of pregnancy	Adjusted effect: 117 g lower than un-exposed. p=0.05	Duration of pregnancy	Family income, maternal height, cigarette smoking, maternal age, parity, child care, antenatal care, child's sex. Not comparable to UK	2+
<i>Cross-sectional</i>						
Saurel-Cubizolles (1987) ¹³⁸	Randomly sampled women who delivered in French maternity units in 1981; n=5508 interviewed, analysis based upon n=2387	Higher rate of low birth weight among women who had to make considerable physical effort, and women who cumulated >2 strenuous working conditions	3.6% with birth weight <2500 g p=0.05	The proportion of women who did not work in the third trimester was higher when the work was strenuous (standing, heavy load carrying, assembly line work, considerable physical effort, p=0.001)	Parity, smoking, preterm delivery	2+
Standing						
<i>Prospective Cohort</i>						
Ha (2002) ⁵⁸	Eligible women who were current employees from 1996–1998 in a petrochemical corporation in Beijing, n=1222 number used in analysis	Standing for >3 hours/day is significantly associated with reduced birth weight	p=0.01	Not noted	Infant gender, maternal age, maternal BMI, maternal education, paternal education, maternal organic solvent exposure	2+
Teitelman (1990) ¹⁵⁹	Eligible women who sought care from various healthcare providers in New Haven, expected to deliver at Yale-New Haven hospital and were pregnant, between 1980 and 1982; n=6219 introduced to the study; n=4186 completed valid interviews; final analysis based upon n=1,206	Mean birth weight lower in those standing, but not significantly. Low birth weight rate was moderately higher, but not statistically significant in those standing	p=0.182 RR 1.40 95% CI 0.64–3.03	Not noted	Parity, smoking, education, caffeine use, illicit drug use, gestational age at interview and marital status	2+

continued

Author	Study population	Main results	Effect measure	Timing in pregnancy	Confounders	Sign grade
Standing and walking						
<i>Prospective cohort</i>						
Henriksen (1995) ⁶⁵	Women attending for antenatal care at Aarhus university hospital from 1989–91, analysis based upon n=4259	Significantly greater risk of preterm delivery if walking and standing for >5 hours/day (does not distinguish between the two), compared with 2 hours or less	Adjusted RR 3.3 95% CI 1.4–8.0	Majority still working after 30 weeks, but a significant number either were off sick or no longer employed. More women who reported jobs involving long hours of standing and walking at 16 weeks, were no longer employed by the 30 th week	Parity, maternal height, smoking, leisure time activities, partner's social class	2+
Working hours						
<i>Prospective cohort</i>						
Tuntiseranee (1998) ¹⁶¹	Pregnant women attending routine antenatal care at the dept of Obstetrics in 2 Thai hospitals from 1994–5, n=1821 number analysed in study n=2084 initially recruited	Significantly increased risk of low birth weight if working 51–60 hours (note unusual trend for >61 hours)	Adjusted OR 2.4 95% CI 1.3–4.4	Not noted	Parity, maternal height, maternal age, obstetrical complications, baby sex, number of antenatal visits, maternal weight at delivery. Not necessarily applicable to UK context	2+
<i>Retrospective cohort</i>						
McDonald (1988) ¹⁰⁸	Women attending 11 obstetrical units in Montreal from 1982–4, n=56,067 interviewed for the study. Results based upon n=22,761	Significant association between working >46 hours/week and low birth weight	O/E 1.34 p<0.01	Not noted	Age, gravidity, previous spontaneous miscarriage, ethnic group, height, educational level, cigarette smoking and alcohol consumption	2++

continued

Physical and shift work in pregnancy: occupational aspects of management

Author	Study population	Main results	Effect measure	Timing in pregnancy	Confounders	Sign grade
Shift work						
<i>Retrospective cohort</i>						
Armstrong (1989) ¹⁴	Women attending 11 obstetrical units in Montreal from 1982–4, n=22,404 number in analysis Further analysis from a previous study	Changing shift work significantly associated with low birth weight before and after accounting for gestational age	Mean per cent predicted birth weight for age 98.2 % 90% CI: 97.2–99.2 p<0.01	Not noted	In previous paper, and then adjusted for occupational sector	2+
Bodin (1999) ²⁵	Female members of the Swedish Midwives Association in 1989; pregnancies between 1980–7, n=1244 number included in analysis	Significant association between night work and preterm birth	Adjusted OR 5.6 95% CI 1.9–16.4	Not noted	Maternal age, parity, employment, work schedule, N2O use	2+
Xu (1994) ¹⁶⁹	Married women working in textile mills in China. Analysis based upon n=887 live births Married, educated, non-smokers, non-alcohol drinkers.	Significant increase in prevalence of low birth weight of all live births associated with rotating shifts Significant association of low birth weight for first live births only and rotating shift work	Adjusted OR 2.1 95% CI 1.1–4.1 p=0.03 Adjusted OR 2.5 95% CI 1.2–5.4 p=0.02	Not noted	Mill location, maternal age at pregnancy, job title, time and duration of leave from the job since pregnancy, order of live births, exposure to dust/gases/fumes, physical activity and position while working, stress at work and use of coal stove for heating at home	2+
<i>Cross-sectional</i>						
Axelsson (1989) ¹⁷	Women who had worked in a particular hospital in Sweden between 1980 and 1984 (previous pregnancies included). Those working nights permanently were selected and a random selection of those not working nights. n=807 sent the questionnaire, analysis based upon n=948 pregnancies	Significantly lower birth weight of second child in non-smoking women working irregular hours, always evenings and rotating shifts.	Irregular hours (p<0.01), always evenings (p<0.01) and rotating shifts (p<0.05)	In the second and third trimester	Not clear, although some results given are 'adjusted'	2+

OR = odds ratio; RR = relative risk

Evidence table 5 Pre-eclampsia

Author	Study population	Main results	Effect measure	Timing in pregnancy	Confounders	Sign grade
Lifting						
<i>Systematic review</i>						
Bonzini (2007) ²⁷	Two studies considered the association between lifting and pre-eclampsia	One study found a positive association between lifting heavy loads and pre-eclampsia	See references in paper	See references in paper	See references in paper	1++
<i>Retrospective cohort</i>						
Wergeland (1997) ¹⁶⁵	Women who gave birth in a 6-week period in 1989 in Norway; singleton pregnancies, n=5438. Analysis based upon n=5388. 3321 in paid work beyond 3rd month of pregnancy	Significant association with lifting heavy loads (10–20 kg) in early pregnancy, compared with not lifting and pre-eclampsia. Significant association with lifting heavy loads >20 times weekly in early pregnancy and pre-eclampsia	Adjusted OR 1.7 95% CI 1.2–2.5 p<0.01 Adjusted OR 2.0 95% CI 1.2–3.2	Early pregnancy	Age, parity, height, BMI, education, smoking, coffee consumption, paid work	2+
Manual handling						
<i>Retrospective cohort</i>						
Wergeland (1997) ¹⁶⁵	Women who gave birth in a 6-week period in 1989 in Norway; singleton pregnancies, n=5438. Analysis based upon n=5388. 3321 in paid work beyond 3rd month of pregnancy	Significant association with work with hands above shoulder level for more than half the day in early pregnancy and pre-eclampsia	Adjusted OR 2.3 95% CI 1.4–3.7	Early pregnancy	Age, parity, height, BMI, education, smoking, coffee consumption, paid work	2+
Working hours						
<i>Systematic review</i>						
Bonzini (2007) ²⁷	Two papers studied the association between working hours and pre-eclampsia	No results given	See references in paper	See references in paper	See references in paper	1++

continued

Physical and shift work in pregnancy: occupational aspects of management

Author	Study population	Main results	Effect measure	Timing in pregnancy	Confounders	Sign grade
Working hours – continued						
<i>Retrospective cohort</i>						
Klebanoff (1990) ⁸¹	Women who graduated from medical school in 1985 and a random sample of male physician spouses Final study population: 989 residents, 1239 spouses. Analysis based upon these numbers.	Residents worked 64 hours/week compared with <37 hours/week in spouses. Twice as many women residents developing pre-eclampsia or eclampsia, compared with spouses	p<0.001	Not noted	Parity, age at delivery, height, pre-pregnancy weight and race or ethnic group	2+
Shift work						
<i>Systematic review</i>						
Bonzini (2007) ²⁷	Two papers studied the association between shift work and pre-eclampsia	No results given	See references in paper	See references in paper	See references in paper	1++
<i>Retrospective cohort</i>						
Wergeland (1997) ¹⁶⁵	Women who gave birth in a 6-week period in 1989 in Norway; singleton pregnancies, n=5438. Analysis based upon n=5388 3321 in paid work beyond 3rd month of pregnancy.	Increased prevalence of pre-eclampsia in parous women doing shift work, however, adjusted OR not significant	Adjusted OR 1.3 95% CI 0.8–1.9 Note: OR 2.0 95% CI 1.1–3.6 for parous women working shifts, compared to women with other schedules.	Work beyond the 3rd month of pregnancy	Age, parity, height, BMI, education, smoking, coffee consumption, paid work	2+
Standing						
<i>Systematic review</i>						
Bonzini (2007) ²⁷	Four papers looked at the association between standing and pre-eclampsia	No results given	See references in paper	See references in paper	See references in paper	1++

OR = odds ratio; RR = relative risk

Evidence table 6 Small for gestational age/ intra-uterine growth restriction

Author	Study population	Main results	Effect measure	Timing in pregnancy	Confounders	Sign grade
Lifting						
<i>Systematic review</i>						
Bonzini (2007) ²⁷	5 studies considered SGA and lifting	No significant positive or negative findings	NA	See references in paper	See references in paper	1++
Physical activity						
<i>Case-control</i>						
Spinillo (1996) ¹⁵²	Cases: Live born singleton infants with a diagnosis of fetal growth restriction delivered in the institution in the study period (1989–94) n=349 Controls: two nulliparous women with appropriate fetal growth during pregnancy, singleton infant born immediately after each case n=698	Increased risk of fetal growth restriction if performing moderate to heavy physical work, compared with light work. Intensity of work based upon type of physical effort (carrying heavy loads, performing heavy cleaning tasks, sweating while on physical exertion). Manual workers and healthcare workers have a significantly higher risk of IUGR	Adjusted OR 2.4 95% CI 1.36–4.21 Adjusted OR 1.81, 1.73 respectively 95% CI 1.15–2.85; 1.06–2.81 respectively	Not noted	Partner's social class, time of stopping work, smoking, alcohol use, illicit drug use, hypertension, maternal age, education, pre-pregnancy BMI, maternal weight gain per week	2+
Standing						
<i>Systematic review</i>						
Bonzini (2007) ²⁷	Association between standing and SGA considered in 8 studies	Majority of risk estimates were <1.5	NA	See references in paper	See references in paper	1++
<i>Prospective cohort</i>						
Teitleman (1990) ¹⁵⁹	Eligible women who sought care from various healthcare providers in New Haven, expected to deliver at Yale-New Haven hospital and were pregnant, between 1980 and 1982; n=6219 introduced to the study n=4186 completed valid interviews; final analysis based upon n=1,206	Rohrer's Index [(birth weight/length ³) x 100] for fetal growth was lower for those with standing jobs, compared with sedentary jobs, however this was not significant	p=0.27	Not noted	Parity, smoking, education, caffeine use, illicit drug use, gestational age at interview and marital status	2+

continued

Physical and shift work in pregnancy: occupational aspects of management

Author	Study population	Main results	Effect measure	Timing in pregnancy	Confounders	Sign grade
Standing – continued						
<i>Retrospective cohort</i>						
Fortier (1995) ⁴⁹	Women in 6 community health departments in Quebec city who gave birth to a live singleton in 1989. Analysis based upon n=4390	Significant risk of SGA baby if standing ≥ 6 hours/day. Risk increases with increasing amount of standing. Significant risk of SGA if works at least 24 weeks and stood for ≥ 6 hours/day	Adjusted OR 1.42 CI 1.02–1.95 Significance of trend p=0.04	Not noted	Age, parity, history of low birth weight or preterm delivery, education, family income, pre-pregnancy weight, height, alcohol and caffeine intake, active and passive smoking, and energy expenditure in non-occupational activities	2++
Working hours						
<i>Systematic review</i>						
Bonzini (2007) ²⁷	Association between working hours and SGA considered in 7 studies	The largest study found an OR of 1.59. 95% CI 1.14–2.22. Most of the rest were close to unity	NA	See references in paper	See references in paper	1++
<i>Prospective cohort</i>						
Tuntiseranee (1998) ¹⁶¹	Pregnant women attending routine antenatal care at the dept of Obstetrics in 2 Thai hospitals from 1994–5, n=1821 number analysed in study; n=2084 initially recruited	Significantly increased risk of small-for-gestational-age if working 51–60 hours (note unusual trend for >61 hours)	Adjusted OR 8.7 95% CI 3.1–24.2	Increased risk of SGA if at 32 weeks continued working >40 hours/week. However small numbers and wide confidence interval	Parity, maternal height, maternal age, obstetrical complications, baby sex, number of antenatal visits, maternal weight at delivery. Not necessarily applicable to UK context	2+
Shift work						
<i>Systematic review</i>						
Bonzini (2007) ²⁷	Association between shift work and SGA considered in 6 studies	In one the RR was 1.5 95% CI 1.0–2.4	Pooled estimate from 6 studies was 1.07 95% CI 0.96–1.19	See references in paper	See references in paper	1++

OR = odds ratio; RR = relative risk

Evidence table 7 Sick leave/sick certification

Author	Study population	Main results	Effect measure	Timing in pregnancy	Confounders	Sign grade
Lifting						
<i>Retrospective cohort</i>						
Kaerlev (2004) ⁷⁷	Women employees aged 20–45 years, in a hospital from 1995–1999 who had a pregnancy. Analysis based upon n=655	Lifting heavy weights significantly associated with >10% sick leave of scheduled working time	Adjusted OR 1.9 CI 1.3–2.9	Lower absence at beginning of pregnancy – increased with gestation. If there was a need for change in work tasks, more sick leave	Age, occupation, full-time or part-time work and previous sickness absence	2+
<i>Cross-sectional</i>						
Strand (1997) ¹⁵⁵	Selection from a population of women in Norway who stayed in the same job during pregnancy and for whom data on leaving work was known. Analysis based upon n=2,713	Significant number left work >3 weeks before delivery if lifting heavy loads (10–20 kg)	Adjusted OR 1.48 CI 1.22–1.80	Not noted	Age, education, children <16 years, housework, sick listed in the year prior to pregnancy	2+
Manual handling						
<i>Retrospective cohort</i>						
Kaerlev (2004) ⁷⁷	Women employees aged 20–45 years, in a hospital from 1995–9 who had a pregnancy. Analysis based upon n=655	Physically demanding jobs significantly associated with sick leave, eg launderers, nursing aides or hospital orderlies and nurses	Not clear if OR are adjusted	Lower absence at beginning of pregnancy – increased with gestation. If there was a need for change in work tasks, more sick leave	Age, occupation, full-time or part-time work and previous sickness absence	2+

continued

Physical and shift work in pregnancy: occupational aspects of management

Author	Study population	Main results	Effect measure	Timing in pregnancy	Confounders	Sign grade
Manual handling – continued						
<i>Cross-sectional</i>						
Strand (1997) ¹⁵⁵	Selection from a population of women in Norway who stayed in the same job during pregnancy and for whom data on leaving work was known. Analysis based upon n=2,713	Significant number left work >3 weeks before delivery if twisting and bending, or working with hands above shoulders; significant number left >8 weeks before delivery if twisting and bending, or working with hands above shoulders. Sick certification increased with increasing level of strain	LSC >3 weeks T + B: Adjusted OR 1.46 CI 1.21–1.76 Hands above shoulder: Adjusted OR: 1.55 CI 1.22–1.95 LSC >8 weeks: T + B: Adjusted OR 1.32 CI 1.05–1.66 Hands above shoulders: Adjusted OR 1.36 CI 1.06–1.74	Not noted	Age, education, children <16 years, housework, sick listed in the year prior to pregnancy	2+
<i>Cross-sectional study taken from a national survey</i>						
Saurel-Cubizolles (1987) ¹³⁸	Randomly sampled women who delivered in French maternity units in 1981. n=5508 interviewed, analysis based upon n=2387	The proportion of women who did not work in the third trimester was higher when the work was strenuous (standing, heavy load carrying, assembly line work, considerable physical effort). More frequent sick leave if work was strenuous. Also more likely to change jobs	p=0.001 p=0.01 and p=0.001	Greater proportion took sick leave or did not work in the 3rd trimester if working conditions were tiring. If working conditions were not modified, there was a higher rate of sick leave, or woman more likely to stop working	Parity, smoking, preterm delivery	2+
Standing						
<i>Cross-sectional</i>						
Strand (1997) ¹⁵⁵	A selection from a population of women in Norway who stayed in the same job during pregnancy and for whom data on leaving work was known. Analysis based upon n=2,713	Significant number left >8 weeks before delivery if standing with their back bent forwards	Adjusted OR 1.30 CI 1.02–1.65	Not noted	Age, education, children <16 years, housework, sick listed in the year prior to pregnancy	2+

continued

Author	Study population	Main results	Effect measure	Timing in pregnancy	Confounders	Sign grade
Standing and walking						
<i>Retrospective cohort</i>						
Kaerlev (2004) ⁷⁷	Women employees aged 20–45 years, in a hospital from 1995–9 who had a pregnancy. Analysis based upon n=655	Standing or walking at work significantly associated with sick leave.	Adjusted OR 3.4 CI 1.9–5.8	Lower absence at beginning of pregnancy – increased with gestation. If there was a need for change in work tasks, more sick leave	Age, occupation, full-time or part-time work and previous sickness absence	2+
Working hours						
<i>Retrospective cohort</i>						
Kaerlev (2004) ⁷⁷	Women employees aged 20–45 years, in a hospital from 1995–9 who had a pregnancy. Analysis based upon n=655	Long working days significantly associated with sick leave	Adjusted OR 3.3 CI 2.2– 4.9	Lower absence at beginning of pregnancy – increased with gestation. If there was a need for change in work tasks, more sick leave	Age, occupation, full-time or part-time work and previous sickness absence	2+
Shift work						
<i>Retrospective cohort</i>						
Kaerlev (2004) ⁷⁷	Women employees aged 20–45 years, in a hospital from 1995–9 who had a pregnancy. Analysis based upon n=655	Night or shift work not significantly associated with sick leave	Adjusted OR 1.4 95 % CI 1.0–1.9	Lower absence at beginning of pregnancy – increased with gestation. If there was a need for change in work tasks, more sick leave	Age, occupation, full-time or part-time work and previous sickness absence	2+
<i>Cross-sectional</i>						
Strand (1997) ¹⁵⁵	Selection from a population of women in Norway who stayed in the same job during pregnancy and for whom data on leaving work was known. Analysis based upon n=2,713	Significant number left work >3 weeks before delivery if doing shift work	Adjusted OR 1.51 CI 1.19–1.93	Not noted	Age, education, children <16 years, housework, sick listed in the year prior to pregnancy	2+

OR = odds ratio; RR = relative risk

Appendix 1 Search strategy

June 1980 – present

1. (pregnant or pregnancy) and work*
2. (pregnant or pregnancy) and employ*
3. (pregnant or pregnancy) and occupation*
4. (pregnant or pregnancy) and manual
5. (expect* mother*) and work*
6. (expect* mother*) and employ*
7. (expect* mother*) and occupation*
8. (expect* mother*) and manual
9. (childbirth) and work*
10. (childbirth) and employ*
11. (childbirth) and occupation*
12. (childbirth) and manual

The above terms were searched with the following words/phrases:

stand*
prolonged standing
morbid*
hypertension
blood pressure
diabet*
premature
preterm
sick*
absenc*
abort*
spontaneous abortion
miscarriage
birth weight
growth retard*
shift*
hour*
excessive hours
nightshift
long hours/long day
bleed*
pain/abdominal
lower back
risk factors

1975 – present

rupture of membranes/preterm rupture of membranes
thrombo-embolic disease/thrombo-embolic disease

Databases and journals searched

BNI
CINAHL
Cochrane Library
Embase
Evidence Based Periodicals Database (*BMJ*)
Health Periodicals Database
HSE Line
Medline
New England Journal of Medicine
Psychinfo
Scisearch

Appendix 2 Critical appraisal form

Reviewer(s): _____

Author, title: _____

Study type (tick all that apply)

- Randomised controlled trial
- Systematic review
- Meta-analysis
- Qualitative research
- Literature review
- Case-control study
- Longitudinal/cohort study
- Other

(Please describe)

Initial comments: _____

SCREENING QUESTIONS

1. Does the paper have a clearly focused aim or research question?

Yes No Can't tell

Consider:

1. population studied
2. interventions delivered
3. outcomes

2. Is the chosen method appropriate?

Yes No Can't tell

Consider whether:

1. the authors explain their research design
2. the chosen method address the research question

Is it worth continuing?

Yes No

Please explain _____

Detailed questions

3. Has the research been conducted rigorously?

Yes No Can't tell

Consider:

1. search strategy described
2. inclusions and exclusions
3. more than one researcher
4. resolving issues of bias

4. Is it clear how data has been analysed?

Yes No Can't tell

Consider:

1. were study results combined
2. if so was this reasonable
3. in-depth description of the analysis process
4. all participants accounted for
5. contradictory findings explained

5. Is there a clear statement of findings?

Yes No Can't tell

Consider:

1. sufficient evidence to support conclusions
2. do findings support the research question
3. precision of results
4. all important variables considered

6. How are the results presented?

Consider:

1. are the results presented numerically, i.e. p-value, OR (odds ratio)
2. are the results presented narratively

7. What is the main result?

Consider:

1. how large is the size of the result
2. how meaningful is the result
3. how would you sum up the bottom-line result in one sentence

8. Are there limitations to the research?

Yes No Can't tell

Consider:

1. was the sample size large enough
2. were all important outcomes considered
3. was the intervention process adequately described
4. was there any follow-up data
5. do the authors acknowledge weaknesses

9. Can the results be applied to a UK context?

Yes No Can't tell

Consider:

1. any discussion on how the findings can be used
2. findings considered in relation to current practice
3. estimation of benefits and costs

Accept for inclusion as evidence Yes No Can't tell

Refer to guideline leader Yes No

Guideline leader's notes

Any references to be followed up from this article?

Please attach this form to your recording sheet for appraising and grading and return to guideline leader

Appendix 3 SIGN grading

Levels of evidence	
1++	High quality meta-analyses, systematic reviews of randomised controlled trials (RCTs) , or RCTs with a very low risk of bias
1+	Well-conducted meta-analyses, systematic reviews of RCTs, or RCTs with a low risk of bias
1–	Meta-analyses, systematic reviews of RCTs, or RCTs with a high risk of bias
2++	High quality systematic reviews of case-control or cohort studies High quality case-control or cohort studies with a very low risk of confounding, bias, or chance and a high probability that the relationship is causal
2+	Well conducted case-control or cohort studies with a low risk of confounding, bias, or chance and a moderate probability that the relationship is causal
2–	Case-control or cohort studies with a high risk of confounding, bias, or chance and a significant risk that the relationship is not causal
3	Non-analytic studies, eg case reports, case series
4	Expert opinion

Appendix 4 Grading system for recommendations

Grades of recommendation	
A	At least one meta-analysis, systematic review, or RCT rated as 1++, and directly applicable to the target population; <i>or</i> A systematic review of RCTs or a body of evidence consisting principally of studies rated as 1+, directly applicable to the target population, and demonstrating overall consistency of results
B	A body of evidence including studies rated as 2++, directly applicable to the target population, and demonstrating overall consistency of results; <i>or</i> Extrapolated evidence from studies rated as 1++ or 1+
C	A body of evidence including studies rated as 2+, directly applicable to the target population and demonstrating overall consistency of results; <i>or</i> Extrapolated evidence from studies rated as 2++
D	Evidence level 3 or 4; <i>or</i> Extrapolated evidence from studies rated as 2+

Good Practice Points

Guideline development groups may find that there is an important practical point that they wish to emphasise but for which there is not, nor is there likely to be, any research evidence. For example, some aspect of management or treatment that is regarded as such sound clinical practice that nobody is likely to question it. These may be marked in the guideline as Good Practice Points. These will not be alternatives to evidence-based recommendations, and will only be used where there is no other way of highlighting the issue.

Appendix 5 Score per paper

Journal: British Journal of Obstetrics and Gynaecology

Title: Standing at work and preterm delivery⁶⁵

Authors: Henriksen TB, Hedegaard M, Secher NJ, Wilcox AJ.

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	✓
	Mostly clear, some information unclear/missing	1	
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	✓
	>400	2	
	100–399	1	
	<100	0	
Response rate	>85%	3	✓
	75–84%	2	
	50–74%	1	
	<50% or unclear	0	
Exposure assessment	Questionnaire: QBasic, Q16, Q30. All completed at home		
Outcome assessment	Ultrasound (82%), LMP or gestational age (9%), birth certificate (9%)		
Inflationary bias	Low	2	✓
	Possible	1	
Confounding	Most confounders considered	2	
	A few confounders considered	1	✓
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	✓
	no	0	
Total			12
SIGN grade			2+

Journal: Occupational & Environmental Medicine

Title: Physical work load and gestational age at delivery⁸³

Authors: Koemeester AP, Broersen JP, Treffers PE.

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	✓
	Mostly clear, some information unclear/missing	1	
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	
	>400	2	
	100–399	1	✓
	<100	0	
Response rate	>85%	3	✓
	75–84%	2	
	50–74%	1	
	<50% or unclear	0	
Exposure assessment	Questionnaire @ 15 weeks. Interview + Questionnaire @ 20 wks		
Outcome assessment	Notes, US (80%) – LMP (20%)		
Inflationary bias	Low	2	✓
	Possible	1	
Confounding	Most confounders considered	2	✓
	A few confounders considered	1	
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	✓
	no	0	
Total			11
SIGN grade			2+

Journal: American Journal of Epidemiology

Title: Prematurity and occupational activity during pregnancy⁹⁸

Authors: Mamelle N, Laumon B, Lazar P.

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	✓
	Mostly clear, some information unclear/missing	1	
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	✓
	>400	2	
	100–399	1	
	<100	0	
Response rate	>85%	3	✓
	75–84%	2	
	50–74%	1	
	<50% or unclear	0	
Exposure assessment	Questionnaire post-delivery		
Outcome assessment	Medicine notes, LMP and Dubowitz score		
Inflationary bias	Low	2	
	Possible	1	✓
Confounding	Most confounders considered	2	✓
	A few confounders considered	1	
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	✓
	no	0	
Total			12
SIGN grade			2+

Journal: International Journal of Epidemiology

Title: Heavy lifting during pregnancy – a hazard to the fetus? A prospective study⁹

Authors: Ahlborg G Jr, Bodin L, Hogstedt C.

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	
	Mostly clear, some information unclear/missing	1	✓
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	✓
	>400	2	
	100–399	1	
	<100	0	
Response rate	>85%	3	✓
	75–84%	2	
	50–74%	1	
	<50% or unclear	0	
Exposure assessment	Questionnaire in early pregnancy prospective, interview post delivery		
Outcome assessment	Delivery records		
Inflationary bias	Low	2	✓
	Possible	1	
Confounding	Most confounders considered	2	
	A few confounders considered	1	✓
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	✓
	no	0	
Total			11
SIGN grade			2+

Journal: Acta Obstetrica et Gynecologia Scandinavia

Title: Standing and walking at work and birthweight⁶⁴

Authors: Henriksen TB, Hedegaard M, Secher NJ.

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	✓
	Mostly clear, some information unclear/missing	1	
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	✓
	>400	2	
	100–399	1	
	<100	0	
Response rate	>85%	3	✓
	75–84%	2	
	50–74%	1	
	<50% or unclear	0	
Exposure assessment	QBasic, Q16 at 16 weeks, and Q30 at 30 weeks		
Outcome assessment	Midwife + Hospital Records		
Inflationary bias	Low	2	✓
	Possible	1	
Confounding	Most confounders considered	2	
	A few confounders considered	1	✓
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	✓
	no	0	
Total			12
SIGN grade			2+

Journal: Epidemiology

Title: Physical exertion as a risk factor for spontaneous abortion⁴²

Authors: Eskenazi B, Fenster L, Wight S, English P, *et al.*

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	✓
	Mostly clear, some information unclear/missing	1	
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	✓
	>400	2	
	100–399	1	
	<100	0	
Response rate	>85%	3	✓
	75–84%	2	
	50–74%	1	
	<50% or unclear	0	
Exposure assessment	Questionnaire months post delivery Computer assisted on telephone		
Outcome assessment	Cases: ?pathology specimen received in path lab. Controls: birth certificates		
Inflationary bias	Low	2	
	Possible	1	✓
Confounding	Most confounders considered	2	✓
	A few confounders considered	1	
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	✓
	no	0	
Total			12
SIGN grade			2+

Journal: Epidemiology**Title:** Pregnancy loss and work schedule during pregnancy⁷⁴**Authors:** Infante-Rivard C, David M, Gauthier R, Rivard GE.

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	✓
	Mostly clear, some information unclear/missing	1	
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	✓
	>400	2	
	100–399	1	
	<100	0	
Response rate	>85%	3	✓
	75–84%	2	
	50–74%	1	
	<50% or unclear	0	
Exposure assessment	Face to face interview in hospital		
Outcome assessment	Not clear in this paper, but does refer reader to another paper for more details		
Inflationary bias	Low	2	
	Possible	1	✓
Confounding	Most confounders considered	2	✓
	A few confounders considered	1	
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	✓
	no	0	
Total			12
SIGN grade			2+

Journal: Epidemiology**Title:** A prospective study of work-related physical exertion and spontaneous abortion⁴⁶**Authors:** Fenster L, Hubbard AE, Windham GC, Waller KO, Swan SH.

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	✓
	Mostly clear, some information unclear/missing	1	
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	✓
	>400	2	
	100–399	1	
	<100	0	
Response rate	>85%	3	✓
	75–84%	2	
	50–74%	1	
	<50% or unclear	0	
Exposure assessment	Interview 1 st trimester – computer assessment telephone interview		
Outcome assessment	Computerised hospital records, medical records/phone some follow-up interviews		
Inflationary bias	Low	2	✓
	Possible	1	
Confounding	Most confounders considered	2	✓
	A few confounders considered	1	
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	✓
	no	0	
Total			13
SIGN grade			2+

Journal: British Journal of Industrial Medicine

Title: Outcome of pregnancy in relation to irregular and inconvenient work schedules¹⁷

Authors: Axelsson G, Rylander R, Molin I.

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	✓
	Mostly clear, some information unclear/missing	1	
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	
	>400	2	✓
	100–399	1	
	<100	0	
Response rate	>85%	3	
	75–84%	2	✓
	50–74%	1	
	<50% or unclear	0	
Exposure assessment	Mailed questionnaire		
Outcome assessment	Mailed questionnaire + verification from hospital records		
Inflationary bias	Low	2	
	Possible	1	✓
Confounding	Most confounders considered	2	
	A few confounders considered	1	✓
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	✓
	no	0	
Total			9
SIGN grade			2+

Journal: Journal of Occupational Health

Title: Prolonged standing and physical exertion at work during pregnancy increases the risk of preterm birth for Thai mothers¹³³

Authors: Ritsmitchai S, Geater AF, Chonsuwiwatvong V.

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	✓
	Mostly clear, some information unclear/missing	1	
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	
	>400	2	✓
	100–399	1	
	<100	0	
Response rate	>85%	3	✓
	75–84%	2	
	50–74%	1	
	<50% or unclear	0	
Exposure assessment	Interviewed by trained interviewers, questionnaires post delivery, for each trimester		
Outcome assessment	Case: Spontaneous preterm abortion,		
	Control: birth at ≥ 37 weeks		
Inflationary bias	Low	2	
	Possible	1	✓
Confounding	Most confounders considered	2	✓
	A few confounders considered	1	
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	✓
	no	0	
Total			11
SIGN grade			2+

Journal: American Journal of Obstetrics & Gynaecology

Title: The association between occupational factors and preterm birth: a United States nurses' study.

Research Committee of the Association of Women's Health, Obstetric, and Neonatal Nurses⁸⁸

Authors: Luke B, Mamelle N, Keith L, Munoz F, *et al.*

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	✓
	Mostly clear, some information unclear/missing	1	
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	✓
	>400	2	
	100–399	1	
	<100	0	
Response rate	>85%	3	
	75–84%	2	
	50–74%	1	
	<50% or unclear	0	✓
Exposure assessment	Mailed survey; structured questionnaire,		
Outcome assessment	LMP, actual delivery date/EDD		
	<37 weeks – cases		
	>37 weeks – controls		
Inflationary bias	Low	2	
	Possible	1	✓
Confounding	Most confounders considered	2	✓
	A few confounders considered	1	
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	✓
	no	0	
Total	Low score because of low percentage response rate, however, total population response was 7,250		9
SIGN grade			2++

Journal: PhD

Title: Perinatal deaths and maternal occupational risk factors in Peninsula Malaysia¹¹

Author: Ali R.

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	✓
	Mostly clear, some information unclear/missing	1	
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	
	>400	2	
	100–399	1	✓
	<100	0	
Response rate	>85%	3	✓
	75–84%	2	
	50–74%	1	
	<50% or unclear	0	
Exposure assessment			
Outcome assessment			
Inflationary bias	Low	2	
	Possible	1	✓
Confounding	Most confounders considered	2	✓
	A few confounders considered	1	
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	✓
	no	0	
Total			10
SIGN grade			2+

Journal: Occupational & Environmental Medicine

Title: Shift work, nitrous oxide exposure, and spontaneous abortion among Swedish midwives¹⁵

Authors: Axelsson G, Ahlborg G Jr, Bodin L.

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	✓
	Mostly clear, some information unclear/missing	1	
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	✓
	>400	2	
	100–399	1	
	<100	0	
Response rate	>85%	3	
	75–84%	2	✓
	50–74%	1	
	<50% or unclear	0	
Exposure assessment	Questionnaire sent at set point to capture previous 10 years of pregnancies. 1 st , 2 nd , & 3 rd trimester information		
Outcome assessment	Reported by woman that was diagnosed as pregnant, or preceded by the pregnancy test		
Inflationary bias	Low	2	
	Possible	1	✓
Confounding	Most confounders considered	2	
	A few confounders considered	1	✓
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	✓
	no	0	
Total			10
SIGN grade			2+

Journal: Epidemiology

Title: The association of shift work and nitrous oxide exposure in pregnancy with birth weight and gestational age²⁵

Authors: Bodin L, Axelsson G, Ahlborg G Jr.

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	✓
	Mostly clear, some information unclear/missing	1	
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	✓
	>400	2	
	100–399	1	
	<100	0	
Response rate	>85%	3	
	75–84%	2	✓
	50–74%	1	
	<50% or unclear	0	
Exposure assessment	E-mailed questionnaire		
Outcome assessment	Questionnaire + Register		
Inflationary bias	Low	2	
	Possible	1	✓
Confounding	Most confounders considered	2	
	A few confounders considered	1	✓
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	✓
	no	0	
Total			10
SIGN grade			2+

Journal: American Journal of Obstetrics & Gynaecology

Title: Occupational fatigue and preterm premature rupture of membranes. National Institute of Child Health and Human Development Maternal-Fetal Medicine, Units Network¹¹⁶

Authors: Newman RB, Goldenberg RL, Moawad AH, Iams JD, *et al.*

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	✓
	Mostly clear, some information unclear/missing	1	
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	✓
	>400	2	
	100–399	1	
	<100	0	
Response rate	>85%	3	✓
	75–84%	2	
	50–74%	1	
	<50% or unclear	0	
Exposure assessment	Nurse administered questionnaires at enrolment (22–4 weeks). Another questionnaire at 27–31 weeks, but no recheck of fatigue		
Outcome assessment	US/LMP/Algorithm used for gestational age		
Inflationary bias	Low	2	✓
	Possible	1	
Confounding	Most confounders considered	2	✓
	A few confounders considered	1	
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	✓
	no	0	
Total			13
SIGN grade			2+

Journal: British Journal of Industrial Medicine

Title: Prematurity and work in pregnancy¹⁰⁸

Authors: McDonald AD, McDonald JC, Armstrong B, Cherry NM *et al.*

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	✓
	Mostly clear, some information unclear/missing	1	
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	✓
	>400	2	
	100–399	1	
	<100	0	
Response rate	>85%	3	✓
	75–84%	2	
	50–74%	1	
	<50% or unclear	0	
Exposure assessment	Interviews soon after delivery		
Outcome assessment	Hospital records		
Inflationary bias	Low	2	
	Possible	1	✓
Confounding	Most confounders considered	2	✓
	A few confounders considered	1	
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	✓
	no	0	
Total			12
SIGN grade			2++

Journal: British Journal of Industrial Medicine

Title: Fetal death and work in pregnancy¹⁰⁷

Authors: McDonald AD, Armstrong B, Cherry NM, Cote R, *et al.*

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	✓
	Mostly clear, some information unclear/missing	1	
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	✓
	>400	2	
	100–399	1	
	<100	0	
Response rate	>85%	3	✓
	75–84%	2	
	50–74%	1	
	<50% or unclear	0	
Exposure assessment	Interviews immediately after delivery		
Outcome assessment	Women and hospital records		
Inflationary bias	Low	2	
	Possible	1	✓
Confounding	Most confounders considered	2	✓
	A few confounders considered	1	
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	✓
	no	0	
Total			12
SIGN grade			2++

Journal: Journal of Occupational Medicine

Title: Spontaneous abortion and occupation¹⁰⁴

Authors: McDonald AD, Armstrong B, Cherry NM, Delorme C *et al.*

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	✓
	Mostly clear, some information unclear/missing	1	
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	✓
	>400	2	
	100–399	1	
	<100	0	
Response rate	>85%	3	✓
	75–84%	2	
	50–74%	1	
	<50% or unclear	0	
Exposure assessment	Interview post delivery		
Outcome assessment	Outcome assessment + medical records		
Inflationary bias	Low	2	
	Possible	1	✓
Confounding	Most confounders considered	2	✓
	A few confounders considered	1	
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	
	no	0	
Total			12
SIGN grade			2+

Journal: Acta Obstetrics & Gynaecology Scandinavia

Title: The effect of work activity in pregnancy on the risk of fetal growth retardation¹⁵²

Authors: Spinillo A, Capuzzo E, Baltaro F, Piazza G, *et al.*

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	✓
	Mostly clear, some information unclear/missing	1	
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	✓
	>400	2	
	100–399	1	
	<100	0	
Response rate	>85%	3	✓
	75–84%	2	
	50–74%	1	
	<50% or unclear	0	
Exposure assessment	Structured interview at birth		
Outcome assessment	US measurements		
Inflationary bias	Low	2	✓
	Possible	1	
Confounding	Most confounders considered	2	✓
	A few confounders considered	1	
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	
	no	0	
Total			12
SIGN grade			2+

Journal: Occupational & Environmental Medicine

Title: Association of rotating shiftwork with preterm births and low birth weight among never smoking women textile workers in China¹⁶⁹

Authors: Xu X, Ding M, Li B, Christiani DC.

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	
	Mostly clear, some information unclear/missing (<i>refers to another paper</i>)	1	✓
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	✓
	>400	2	
	100–399	1	
	<100	0	
Response rate	>85%	3	✓
	75–84%	2	
	50–74%	1	
	<50% or unclear	0	
Exposure assessment	Standardised questionnaire, but not clear at what point is pregnancy		
Outcome assessment	Not clear, but refers to another paper		
Inflationary bias	Low	2	
	Possible	1	✓
Confounding	Most confounders considered	2	✓
	A few confounders considered	1	
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	✓
	no	0	
Total			11
SIGN grade			2+

Journal: Journal of Epidemiology & Community Health

Title: Employment, working conditions, and preterm birth: results from the Europop case-control survey¹⁴⁰

Authors: Saurel-Cubizolles MJ, Zeitlin J, Lelong N, Papiernik E *et al.*

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	✓
	Mostly clear, some information unclear/missing	1	
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	✓
	>400	2	
	100–399	1	
	<100	0	
Response rate	>85%	3	✓
	75–84%	2	
	50–74%	1	
	<50% or unclear	0	
Exposure assessment	Interviews after delivery		
Outcome assessment	Medical records		
Inflationary bias	Low	2	
	Possible	1	✓
Confounding	All confounders considered	2	✓
	A few confounders considered	1	
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	✓
	no	0	
Total			12
SIGN grade			2++

Journal: American Journal of Obstetrics & Gynaecology

Title: Shift work, duration of pregnancy, and birth weight: the National Birth Cohort in Denmark¹⁷²

Authors: Zhu JL, Hjollund NH, Olsen J.

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	✓
	Mostly clear, some information unclear/missing	1	
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	✓
	>400	2	
	100–399	1	
	<100	0	
Response rate	>85%	3	✓
	75–84%	2	
	50–74%	1	
	<50% or unclear	0	
Exposure assessment	1 st Interviews ≥11–25 weeks		
	2 nd Interviews – 27–37 weeks		
Outcome assessment	LMP at prenatal visit, EDD at 2 nd interviews; US measurement		
Inflationary bias	Low	2	✓
	Possible	1	
Confounding	Most confounders considered	2	✓
	A few confounders considered	1	
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	✓
	no	0	
Total			13
SIGN grade			2++

Journal: The New England Journal of Medicine

Title: Outcomes of pregnancy in a national sample of resident physicians⁸¹

Author: Klebanoff MA, Shiono PH, Rhoads GG.

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	
	Mostly clear, some information unclear/missing	1	✓
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	✓
	>400	2	
	100–399	1	
	<100	0	
Response rate	>85%	3	✓
	75–84%	2	
	50–74%	1	
	<50% or unclear	0	
Exposure assessment	Questionnaire at what point?		
	Implied post delivery		
Outcome assessment	Not clear – but was asked in questionnaire		
Inflationary bias	Low	2	
	Possible	1	✓
Confounding	Most confounders considered	2	
	A few confounders considered	1	✓
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	✓
	no	0	
Total			10
SIGN grade			2+

Journal: American Journal of Epidemiology

Title: Tree based, Two-stage Risk factor analysis for spontaneous abortion¹⁷⁰

Authors: Zhang H, Bracken MB

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	
	Mostly clear, some information unclear/missing	1	✓
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	✓
	>400	2	
	100–399	1	
	<100	0	
Response rate	>85%	3	✓
	75–84%	2	
	50–74%	1	
	<50% or unclear	0	
Exposure assessment	Home interviews 5–16 weeks		
Outcome assessment	Details in: Bracken MB, Belanger K, Hellenbrand K <i>et al.</i> *		
Inflationary bias	Low	2	✓
	Possible	1	
Confounding	Most confounders considered	2	✓
	A few confounders considered	1	
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	✓
	no	0	
Total			12
SIGN grade			2+

*Details in: Bracken MB, Belanger K, Hellenbrand K *et al.* Exposure to electromagnetic fields during pregnancy with emphasis on electrically heated beds: association with birth weight and intra-uterine growth retardation. *Epidemiology* 1995;6:263–70.

Journal: British Journal of Industrial Medicine

Title: Work in pregnancy and birth weight for gestational age¹⁴

Authors: Armstrong BG, Nolin AD, McDonald AD.

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	
	Mostly clear, some information unclear/missing	1	✓
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	✓
	>400	2	
	100–399	1	
	<100	0	
Response rate	>85%	3	✓
	75–84%	2	
	50–74%	1	
	<50% or unclear	0	
Exposure assessment	Interview using questionnaire soon after delivery		
Outcome assessment			
Inflationary bias	Low	2	
	Possible	1	✓
Confounding	Most confounders considered	2	✓
	A few confounders considered	1	
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	✓
	no	0	
Total			11
SIGN grade			2+

Journal: Journal of Occupational & Environmental Medicine

Title: Does standing at work during pregnancy result in reduced infant birth weight?⁵⁸

Authors: Ha E, Cho SI, Park H, Chen D *et al.*

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	
	Mostly clear, some information unclear/missing	1	✓
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	✓
	>400	2	
	100–399	1	
	<100	0	
Response rate	>85%	3	✓
	75–84%	2	
	50–74%	1	
	<50% or unclear	0	
Exposure assessment	Administered questionnaire but not clear when		
Outcome assessment	Hospital delivery records		
Inflationary bias	Low	2	
	Possible	1	✓
Confounding	Most confounders considered	2	✓
	A few confounders considered	1	
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	✓
	no	0	
Total			11
SIGN grade			2+

Journal: Journal of Occupational and Environmental Medicine

Title: Shift work, job stress, and late fetal loss: The National Birth Cohort in Denmark¹⁷¹

Authors: Zhu JL, Hjollund NH, Andersen AM, Olsen, J.

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	✓
	Mostly clear, some information unclear/missing	1	
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	✓
	>400	2	
	100–399	1	
	<100	0	
Response rate	>85%	3	✓
	75–84%	2	
	50–74%	1	
	<50% or unclear	0	
Exposure assessment	1 st interviews 11–25 weeks		
	2 nd interviews		
Outcome assessment	LMP, EDD and register		
Inflationary bias	Low	2	✓
	Possible	1	
Confounding	Most confounders considered	2	✓
	A few confounders considered	1	
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	✓
	no	0	
Total			13
SIGN grade			2+

Journal: Scandinavian Journal of Public Health

Title: Long-term sick leave and its risk factors during pregnancy among Danish hospital employees⁷⁷

Authors: Kaerlev L, Jacobsen LB, Olsen J, Bonde JP.

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	✓
	Mostly clear, some information unclear/missing	1	
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	
	>400	2	✓
	100–399	1	
	<100	0	
Response rate	>85%	3	✓
	75–84%	2	
	50–74%	1	
	<50% or unclear	0	
Exposure assessment	From hospital roster, questionnaire considered changes during pregnancy		
Outcome assessment	Register date, roster		
Inflationary bias	Low	2	
	Possible	1	✓
Confounding	Most confounders considered	2	
	A few confounders considered	1	✓
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	✓
	no	0	
Total			10
SIGN grade			2+

Physical and shift work in pregnancy: occupational aspects of management

Journal: Journal of Epidemiology and Community Health

Title: Is preterm delivery still related to physical working conditions in pregnancy?¹⁴¹

Authors: Saurel-Cubizolles MJ, Subtil D, Kaminski M.

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	✓
	Mostly clear, some information unclear/missing	1	
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	✓
	>400	2	
	100–399	1	
	<100	0	
Response rate	>85%	3	✓
	75–84%	2	
	50–74%	1	
	<50% or unclear	0	
Exposure assessment	Structured questionnaire after delivery in hospital		
Outcome assessment	LMP, obstetric staff estimates		
Inflationary bias	Low	2	
	Possible	1	✓
Confounding	Most confounders considered	2	
	A few confounders considered	1	✓
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	✓
	no	0	
Total			11
SIGN grade			2+

Journal: Scand Journal of Social Medicine

Title: Work load, job control and risk of leaving work by sickness certification before delivery, Norway 1989¹⁵⁵

Authors: Strand K, Wergeland E, Bjerkedal T.

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	✓
	Mostly clear, some information unclear/missing	1	
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	✓
	>400	2	
	100–399	1	
	<100	0	
Response rate	>85%	3	✓
	75–84%	2	
	50–74%	1	
	<50% or unclear	0	
Exposure assessment	Questionnaire after delivery in hospital		
Outcome assessment	Questionnaire answers + some checking with a database		
Inflationary bias	Low	2	
	Possible	1	✓
Confounding	Most confounders considered	2	✓
	A few confounders considered	1	
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	✓
	no	0	
Total			12
SIGN grade			2+

Journal: American Journal of Epidemiology

Title: Effect of maternal work activity on preterm birth and low birth weight¹⁵⁹

Authors: Teitelman AM, Welch LS, Hellenbrand KG, Bracken MB.

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	✓
	Mostly clear, some information unclear/missing	1	
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	✓
	>400	2	
	100–399	1	
	<100	0	
Response rate	>85%	3	✓
	75–84%	2	
	50–74%	1	
	<50% or unclear	0	
Exposure assessment	Interviewed using a standard questionnaire in 1 st or 2 nd trimester		
Outcome assessment	Hospital records or medical provider		
Inflationary bias	Low	2	✓
	Possible	1	
Confounding	Most confounders considered	2	✓
	A few confounders considered	1	
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	✓
	no	0	
Total			13
SIGN grade			2+

Journal: Journal of Occupational & Environmental Medicine

Title: The effect of heavy maternal workload on fetal growth retardation and preterm delivery. A study among southern Thai women¹⁶¹

Authors: Tuntiseranee P, Geater A, Chongsuivatwong V, Kor-anantakul O.

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	✓
	Mostly clear, some information unclear/missing	1	
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	✓
	>400	2	
	100–399	1	
	<100	0	
Response rate	>85%	3	✓
	75–84%	2	
	50–74%	1	
	<50% or unclear	0	
Exposure assessment	Q17, Q32 – interviews by trained interviewers		
Outcome assessment	From obstetric notes after delivery		
Inflationary bias	Low	2	✓
	Possible	1	
Confounding	Most confounders considered	2	✓
	A few confounders considered	1	
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	✓
	no	0	
Total			13
SIGN grade			2+

Journal: International Journal of Epidemiology

Title: Influence of heavy agricultural work during pregnancy on birthweight in northeast Brazil⁸⁶

Authors: Lima M, Ismail S, Ashworth A, Morris SS.

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	✓
	Mostly clear, some information unclear/missing	1	
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	
	>400	2	✓
	100–399	1	
	<100	0	
Response rate	>85%	3	✓
	75–84%	2	
	50–74%	1	
	<50% or unclear	0	
Exposure assessment	Hospital based interviews 12–48 hours post delivery – trained interviewers		
Outcome assessment	'Antropometric' measures		
Inflationary bias	Low	2	
	Possible	1	✓
Confounding	Most confounders considered	2	✓
	A few confounders considered	1	
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	✓
	no	0	
Total			11
SIGN grade			2+

Journal: British Journal of Industrial Medicine

Title: Pregnant women's working conditions and their changes during pregnancy: a national study in France¹³⁸

Authors: Saurel-Cubizolles MJ, Kaminski M.

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	✓
	Mostly clear, some information unclear/missing	1	
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	✓
	>400	2	
	100–399	1	
	<100	0	
Response rate	>85%	3	✓
	75–84%	2	
	50–74%	1	
	<50% or unclear	0	
Exposure assessment	Interviews within 6 days of delivery, standard questionnaire, 1 st trimester + modifications		
Outcome assessment	Preterm <37 weeks, low <2500g		
Inflationary bias	Low	2	
	Possible	1	✓
Confounding	Most confounders considered	2	
	A few confounders considered	1	✓
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	✓
	no	0	
Total			11
SIGN grade			2+

Journal: International Journal of Gynaecology & Obstetrics

Title: Working conditions and prevalence of pre-eclampsia, Norway 1989¹⁶⁵

Authors: Wergeland E, Strand K.

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	✓
	Mostly clear, some information unclear/missing	1	
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	✓
	>400	2	
	100–399	1	
	<100	0	
Response rate	>85%	3	✓
	75–84%	2	
	50–74%	1	
	<50% or unclear	0	
Exposure assessment	Questionnaires post delivery		
Outcome assessment	Report by women possibly cross-referenced with medical notes, but not clear		
Inflationary bias	Low	2	
	Possible	1	✓
Confounding	Most confounders considered	2	✓
	A few confounders considered	1	
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	✓
	no	0	
Total			12
SIGN grade			2+

Journal: Scandinavian Journal of Work Environmental Health

Title: Maternal work during pregnancy and the risks of delivering a small-for-gestational-age or preterm infant⁴⁹

Authors: Fortier I, Marcoux S, Brisson J.

Points to score	Conditions	Score	Points
Sampling	Defined population, representative, clear procedures, all accounted for	2	
	Mostly clear, some information unclear/missing	1	✓
	Mostly unclear, difficult to track or unmentioned	0	
Power	>1000	3	✓
	>400	2	
	100–399	1	
	<100	0	
Response rate	>85%	3	✓
	75–84%	2	
	50–74%	1	
	<50% or unclear	0	
Exposure assessment	Telephone interviews post delivery		
Outcome assessment	Birth weight from birth certificate.		
	Gestational age from US/LMP/birth certificate		
Inflationary bias	Low	2	
	Possible	1	✓
Confounding	All confounders considered	2	✓
	A few confounders considered	1	
	None considered	0	
Statistics	OR/95%/p-value included: yes	1	✓
	no	0	
Total			11
SIGN grade			2++

References

- 1 The Control of Lead at Work Regulations 2002. SI 2002/2676. Webpage: www.opsi.gov.uk/si/si2002/20022676.htm
- 2 The Ionising Radiations Regulations 1999. SI 1999/3232. Webpage: www.opsi.gov.uk/si/si1999/19993232.htm
- 3 The Management of Health and Safety at Work Regulations 1999. SI 1999/3242. Webpage: www.opsi.gov.uk/si/si1999/19993242.htm
- 4 Employment Rights Act 1996. Webpage: www.opsi.gov.uk/acts/acts1996/Ukpga_19960018_en_1
- 5 Employment Relations Act 1999. Webpage: www.opsi.gov.uk/ACTS/acts1999/pdf/ukpga_19990026_en.pdf
- 6 The Workplace (Health, Safety and Welfare) Regulations 1992. SI 1992/3004. Webpage: www.opsi.gov.uk/SI/si1992/Uksi_19923004_en_1.htm
- 7 Sex Discrimination Act 1975. Webpage: www.opsi.gov.uk/acts/acts1975/PDF/ukpga_19750065_en.pdf
- 8 Ahlborg G Jr. Physical work load and pregnancy outcome. *J Occup Environ Med* 1995 Aug;37(8):941–4.
- 9 Ahlborg G Jr, Bodin L, Hogstedt C. Heavy lifting during pregnancy – a hazard to the foetus? A prospective study. *Int J Epidemiol* 1990 Mar;19(1):90–7.
- 10 Ahlborg G Jr, Hogstedt C, Bodin L, Barany S. Pregnancy outcome among working women. *Scand J Work Environ Health* 1989 Jun;15(3):227–33.
- 11 Ali R. Perinatal deaths and maternal occupational risk factors in Peninsular Malaysia. PhD. 2002.
- 12 News about the effects of work and exercise in pregnancy. *Child Health Alert*. 1990 May;3–4.
- 13 Premature birth linked with prolonged standing. *The Women's Letter*. 1990 Jan;3(1):5.
- 14 Armstrong BG, Nolin AD, McDonald AD. Work in pregnancy and birth weight for gestational age. *Br J Ind Med* 1989 Mar;46(3):196–9.
- 15 Axelsson G, Ahlborg G Jr, Bodin L. Shift work, nitrous oxide exposure, and spontaneous abortion among Swedish midwives. *Occup Environ Med* 1996 Jun;53(6):374–8.
- 16 Axelsson G, Lutz C., Rylander R. Exposure to solvents and outcome of pregnancy in university laboratory employees. *Br J Ind Med* 1984 Aug;41(3):305–12.
- 17 Axelsson G, Rylander R, Molin I. Outcome of pregnancy in relation to irregular and inconvenient work schedules. *Br J Ind Med* 1989 Jun;46(6):393–8.
- 18 Bannerjee B, Dey TK, Chatterjee P. Estimation of risk of pregnancy wastage due to lifting of heavy weight during pregnancy. *Indian Journal of Occupational and Environmental Medicine* Vol.Iss/Pg. 6/1 (13–15).
- 19 BaoYS. The effects of carrying and moving heavy loads on female reproductive system and maternal function. *Zhonghua Yu Fang Yi Xue Za Zhi*. 1989 Jul; 23(4):195–8.
- 20 Barrett V, Phillips JA. Reproductive health in the American workplace. *AAOHN J* 1995 Jan;43(1):40–8; quiz 49–51.
- 21 Bergsjö P, Villar J. Scientific basis for the content of routine antenatal care. II. Power to eliminate or alleviate adverse newborn outcomes; some special conditions and examinations. *Acta Obstet Gynecol Scand*. 1997 Jan; 76(1):15–25.
- 22 Berkowitz GS. Employment-related physical activity and pregnancy outcome. *J Am Med Womens Assoc* 1995 Sep-1995 Oct 31; 50(5):167–9, 174.
- 23 Berkowitz GS, Papiernik E. Working conditions, maternity legislation, and preterm birth. *Semin Perinatol* 1995 Aug;19(4):272–8.
- 24 Bialobok KM, Monga M. Fatigue and work in pregnancy. *Curr Opin Obstet Gynecol* 2000 Dec; 12(6):497–500.
- 25 Bodin L, Axelsson G, Ahlborg G Jr. The association of shift work and nitrous oxide exposure in pregnancy with birth weight and gestational age. *Epidemiology* 1999 Jul;10(4):429–36.
- 26 Bonjour D, Lissenburgh S. Maternity Rights in Britain 2002 : Survey of Employers. A report by the Policy Studies Institute for the DWP. Department of Work & Pensions. 2002;In-house Report 130.

- 27 Bonzini M, Coggon D, Palmer KT. Risk of prematurity, low birthweight and pre-eclampsia in relation to working hours and physical activities: a systematic review. *Occup Environ Med* 2007 Apr;64(4): 228–43.
- 28 Brinville S. The pregnant or employee. Part 1 Ensuring maternal health. *AORN Journal* 1987 Feb; 45(2):pgs 404–410.
- 29 Bryant HE, Love E J. Effect of employment and its correlates on spontaneous abortion risk. *Soc Sci Med* 1991;33(7):795–800.
- 30 Catherine Nelson-Piercy. *Handbook of obstetric medicine*, 2nd edn. Oxford: Taylor & Francis Ltd, 2001:288.
- 31 Ceron-Mireles P, Harlow SD, Sanchez-Carrillo CI. The risk of prematurity and small-for-gestational-age birth in Mexico City: the effects of working conditions and antenatal leave. *Am J Public Health* 1996 Jun; 86(6):825–31.
- 32 Ceron-Mireles P, Harlow SD, Nunex-Urquiza RM. Maternal working conditions and low birth weight in Mexico City. *Salud Publica De Mexico* 1997;39(1):2–10.
- 33 Ceron-Mireles P, Harlow SD. Low birth weight and occupational stress in Mexico City. *Am J Epidemiol* 1994;Jun;vol 139, no 11 suppl S75.
- 34 Chavkin W. Work and pregnancy. Review of the literature and policy discussion. *Obstet Gynecol Surv.* 1986 Aug;41(8):467–72.
- 35 Chien LY, Ko YL. Fatigue during pregnancy predicts caesarean deliveries. *J Adv Nurs* 2004 Mar; 45(5): 487–94.
- 36 Clapp JF. 3rd. Pregnancy outcome: physical activities inside versus outside the workplace. *Semin Perinatol* 1996 Feb;20(1):70–6.
- 37 Colie CF. Preterm labour and delivery in working women. *Semin Perinatol* 1993 Feb;17(1):37–44.
- 38 Conrad W. A study of the management of new and expectant mothers in the National Health Service Occupational Health Practice. MFOM Dissertation. 2000.
- 39 Department of Work & Pensions. *Work & pensions statistics*, 32nd ed. National Statistics, London: DWP, 2004.
- 40 Department of Work & Pensions. *A guide to maternity benefits – April 2006*. London: DWP, 2006: NI 17A
- 41 Eriksen G, Wohlert M, Ersbak V *et al.* Placental abruption. A case-control investigation. *Br J Obstet Gynaecol* 1991 May;98(5):448–52.
- 42 Eskenazi B, Fenster L, Wight S, English P, *et al.* Physical exertion as a risk factor for spontaneous abortion. *Epidemiology* 1994 Jan;5(1):6–13.
- 43 European Union Council. The introduction of measures to encourage improvements in the safety and health at work of pregnant workers and workers who have recently given birth or are breastfeeding. European Union Council Directive . (92/85/EEC).
Webpage: <http://europa.eu/scadplus/leg/en/cha/c10914.htm>
- 44 Fall CHD. Developmental origins of cardiovascular disease, Type 2 diabetes and obesity in humans. In: Wintour-Coghlan M, Owens J (eds), *Early life origins of health and disease*. New York: Landes Bioscience, 2006.
- 45 Feinberg JS, Kelley CR. Pregnant workers. A physician's guide to assessing safe employment. *West J Med* 1998 Feb;168(2):86–92.
- 46 Fenster L, Hubbard AE, Windham GC, Waller KO, Swan SH. A prospective study of work-related physical exertion and spontaneous abortion. *Epidemiology* 1997 Jan;8(1):66–74.
- 47 Florack EI, Pellegrino AE, Zielhuis GA, Rolland R. Influence of occupational physical activity on pregnancy duration and birthweight. *Scand J Work Environ Health* 1995 Jun;21(3):199–207.
- 48 Florack EI, Zielhuis GA, Pellegrino JE, Rolland R. Occupational physical activity and the occurrence of spontaneous abortion. *Int J Epidemiol* 1993 Oct;22(5):878–84.
- 49 Fortier I, Marcoux S, Brisson J. Maternal work during pregnancy and the risks of delivering a small-for-gestational-age or preterm infant. *Scand J Work Environ Health* 1995 Dec;21(6):412–8.
- 50 Frazier LM. Workplace reproductive problems. *Prim Care* 2000 Dec;27(4):1039–56.

- 51 Frazier LM, Golbeck AL, Lipscomb L. Medically recommended cessation of employment among pregnant women in Georgia. *Obstet Gynecol* 2001 Jun;97(6):971–5.
- 52 Frazier LM, Ho HL, Molgaard CA. Variability in physician management of employment during pregnancy. *Women Health* 2001;34(4):51–63.
- 53 Gabbe SG, Morgan MA, Power ML, Schulkin J, Williams SB. Duty hours and pregnancy outcome among residents in obstetrics and gynaecology. *Obstet Gynecol* 2003 Nov;102(5 Pt 1):948–51.
- 54 Gilmour D. Risk for the new or expectant mother working in the perioperative environment. *Br J Perioper Nurs* 2000 Jun;10(6):306–10.
- 55 Gold EB, Tomich E. Occupational hazards to fertility and pregnancy outcome. *Occup Med* 1994 Jul-1994 Sep 30;9(3):435–69.
- 56 Goulet L, Theriault G. Association between spontaneous abortion and ergonomic factors. A literature review of the epidemiologic evidence. *Scand J Work Environ Health* 1987 Oct;13(5):399–403.
- 58 Ha E, Cho SI, Park H, Chen D *et al*. Does standing at work during pregnancy result in reduced infant birth weight? *J Occup Environ Med* 2002 Sep;44(9):815–21.
- 59 Hanke W, Kalinka J, Makowicz-Dabrowska T, Sobala W. Heavy physical work during pregnancy – a risk factor for small-for-gestational-age babies in Poland. *Am J Ind Med* 1999 Jul;36(1):200–5.
- 60 Haram K, Mortensen JH, Wollen AL. Preterm delivery: an overview. *Acta Obstet Gynecol Scand* 2003 Aug;82(8):687–704;ISSN:0001–6349 (Print).
- 61 Gross H, Patterson H. Pregnancy and working: a critical reading of advice and information on pregnancy and employment. *Feminism & Psychology* 2001;11(4):511–525.
- 62 Hatch M, Ji BT, Shu XO, Susser M. Do standing, lifting, climbing, or long hours of work during pregnancy have an effect on foetal growth? *Epidemiology* 1997 Sep;8(5):530–6.
- 63 Hatch M, Moline J. Women, work, and health. *Am J Ind Med* 1997 Sep;32(3):303–8.
- 64 Henriksen TB, Hedegaard M, Secher NJ. Standing and walking at work and birthweight. *Acta Obstet Gynecol Scand* 1995 Aug; 74(7):509–16;ISSN:0001–6349
- 65 Henriksen TB, Hedegaard M, Secher NJ, Wilcox AJ. Standing at work and preterm delivery. *Br J Obstet Gynaecol* 1995 Mar;102(3):198–206.
- 66 Hickey CA, Cliver SP, Mulvihill FX, McNeal SF *et al*. Employment-related stress and preterm delivery: a contextual examination. *Public Health Rep* 1995 Jul-1995 Aug 31;110(4):410–8.
- 67 Higgins JR, Walshe JJ, Conroy RM, Darling MR. The relation between maternal work, ambulatory blood pressure, and pregnancy hypertension. *J Epidemiol Community Health* 2002 May;56(5):389–93.
- 68 Hjollund NH, Jensen TK, Bonde JP, Henriksen TB *et al*. Spontaneous abortion and physical strain around implantation: a follow-up study of first-pregnancy planners. *Epidemiology* 2000 Jan;11(1):18–23.
- 69 Homer CJ, Beresford SA, James SA, Siegel E, Wilcox S. Work-related physical exertion and risk of preterm, low birthweight delivery. *Paediatr Perinat Epidemiol* 1990 Apr;4(2):161–74.
- 70 Hrubá D, Kukla L, Tyrlik M. Occupational risks for human reproduction: ELSPAC Study. European Longitudinal Study of Pregnancy and Childhood. *Cent Eur J Public Health* 1999 Nov;7(4):210–5.
- 71 Health and Safety Executive. *A guide for new and expectant mothers who work*. London: HSE, 2003.
- 72 Health and Safety Executive. *New and expectant mothers at work – a guide for health professionals*. London: HSE, 2003.
- 73 Hudson M, Lissenburgh S, Sohir-Dikmen M. *Maternity & paternity rights in Britain 2002: a survey of parents*. A report by the Policy Studies Institute for the DWP & the DTI. In-house Report 131. London: Department of Work & Pensions, 2002.
- 74 Infante-Rivard C, David M, Gauthier R, Rivard GE. Pregnancy loss and work schedule during pregnancy. *Epidemiology* 1993 Jan;4(1):73–5.
- 75 Irwin DE, Savitz DA, St Andre KA, Hertz-Picciotto I. Study of occupational risk factors for pregnancy-induced hypertension among active duty enlisted Navy personnel. *Am J Ind Med* 1994 Mar;25(3):349–59.
- 76 Johnson D. Stand and deliver. *Vegetarian Times* 2000 Jul;16
- 77 Kaerlev L, Jacobsen LB, Olsen J, Bonde JP. Long-term sick leave and its risk factors during pregnancy among Danish hospital employees. *Scand J Public Health* 2004;32(2):111–7.
- 78 Katz VL, Jenkins T, Haley L, Bowes WA Jr. Catecholamine levels in pregnant physicians and nurses: a pilot study of stress and pregnancy. *Obstet Gynecol* 1991 Mar;77(3):338–42.

- 79 Killien MG. Women and employment: a decade review. *Annu Rev Nurs Res* 2001;19:87–123.
- 80 Klebanoff MA, Shiono PH, Carey JC. The effect of physical activity during pregnancy on preterm delivery and birth weight. *Am J Obstet Gynecol* 1990 Nov;163(5 Pt 1):1450–6.
- 81 Klebanoff MA, Shiono PH, Rhoads GG. Outcomes of pregnancy in a national sample of resident physicians. *N Engl J Med* 1990 Oct 11;323(15):1040–5.
- 82 Knutsson A. Health disorders of shift workers. *Occup Med (Lond)* 2003 Mar;53(2):103–8.
- 83 Koemeester AP, Broersen JP, Treffers PE. Physical work load and gestational age at delivery. *Occup Environ Med* 1995 May;52(5):313–5.
- 84 Kogi K, Miura T. Shiftwork: its practice and improvement. Proceedings of the Sixth International Symposium on Night and Shift Work. Kyoto, 30 August-1 September 1982. *J Hum Ergol (Tokyo)* 1982;11 Suppl:1–541.
- 85 Launer LJ, Villar J, Kestler E, de Onis M. The effect of maternal work on foetal growth and duration of pregnancy: a prospective study. *Br J Obstet Gynaecol* 1990 Jan;97(1):62–70.
- 86 Lima M, Ismail S, Ashworth A, Morris SS. Influence of heavy agricultural work during pregnancy on birthweight in northeast Brazil. *Int J Epidemiol* 1999 Jun;28(3):469–74.
- 87 Lindsay S. The influence of the ‘spaces of everyday life’ on pregnancy health. *The Canadian Geographer*. 2004 Mar;48(1):35–51.
- 88 Luke B, Mamelle N, Keith L, Munoz F, *et al*. The association between occupational factors and preterm birth: a United States nurses’ study. Research Committee of the Association of Women’s Health, Obstetric, and Neonatal Nurses. *Am J Obstet Gynecol* 1995 Sep;173(3 Pt 1):849–62.
- 89 Lyness KS, Thompson CA, Francesco AM, Judiesch MK. Work and pregnancy: individual and organizational factors influencing organizational commitment, timing of maternity leave, and return to work. *Sex Roles* 1999 Oct;41(7-8):485–508.
- 90 Macfarlane AJ, Mugford M. *Birth Counts – statistics of pregnancy and childbirth*, 2nd edn. London: The Stationery Office, 2000.
- 91 MacKey MC, Williams CA, Tiller C. M. Stress, pre-term labour and birth outcomes. *J Adv Nurs* 2000 Sep;32(3):666–74.
- 92 Magann EF, Nolan TE. Pregnancy outcome in an active-duty population. *Obstet Gynecol* 1991 Sep; 78(3 Pt 1):391–3.
- 93 Makoweic-Dabrowska T, Siedlecka J. Occupational physical activity and the course of pregnancy and its outcome. *Medycyna-Pracy* 47(6):629–649.
- 94 Makowiec-Dabrowska T, Hanke W, Radwan-Wlodarczyk Z, Koszada-Wlodarczyk W, Sobala, W. Working condition of pregnant women. Departures from regulation on occupations especially noxious or hazardous to women. *Med Pr* 2003;54(1):33–43.
- 95 Makowiec-Dabrowska T, Hanke W, Sobala W, Radwan-Wlodarczyk Z, Koszada-Wlodarczyk W. Risk of certain obstetric pathologies in women employed in working conditions non-complying with the current legal status on work load and working conditions appropriate for pregnant women. *Med Pr* 2003;54(5):415–25.
- 96 Makowiec-Dabrowska T, Siedlecka J. Physical exertion at work and the course and outcome of pregnancy. *Med Pr* 1996;47(6):629–49.
- 97 Mamelle N, Bertucat I, Munoz F. Pregnant women at work: rest periods to prevent preterm birth? *Paediatr Perinat Epidemiol* 1989 Jan;3(1):19–28.
- 98 Mamelle N, Laumon B, Lazar P. Prematurity and occupational activity during pregnancy. *Am J Epidemiol* 1984 Mar;119(3):309–22.
- 99 Mamelle N, Munoz F. Occupational working conditions and preterm birth: A reliable scoring system. *American Journal of Epidemiology* 1987;Vol.126(1) pp150–152.
- 100 Marbury MC. Adverse working conditions and premature delivery. *Am J Public Health* 1991 Aug; 81(8):973–4.
- 101 Marcoux S, Berube S, Brisson C, Mondor M. Job strain and pregnancy-induced hypertension. *Epidemiology* 1999 Jul;10(4):376–82.
- 102 McCulloch J. Health risks associated with prolonged standing. *Work* 2002;19(2):201–5.
- 103 McDonald AD. Work and pregnancy. *Br J Ind Med* 1988 Sep;45(9):577–80.

- 104 McDonald AD, Armstrong B, Cherry NM, Delorme C *et al.* Spontaneous abortion and occupation. *J Occup Med* 1986 Dec;28(12):1232–8.
- 105 McDonald AD, McDonald JC, Armstrong B, Cherry N *et al.* Occupation and pregnancy outcome. *Br J Ind Med* 1987 Aug;44(8):521–6.
- 106 McDonald AD, McDonald JC, Armstrong B, Cherry N *et al.* Congenital defects and work in pregnancy. *Br J Ind Med* 1988 Sep;45(9):581–8.
- 107 McDonald AD. Foetal death and work in pregnancy. *Br J Ind Med* 1988 Mar;45(3):148–57.
- 108 McDonald AD, McDonald JC, Armstrong B, Cherry NM *et al.* Prematurity and work in pregnancy. *Br J Ind Med* 1988 Jan;45(1):56–62.
- 109 McGill M. Work and Pensions Statistics 2004;1–251.
- 110 Mercer BM, Goldenberg RL, Das A, Moawad AH *et al.* The preterm prediction study: a clinical risk assessment system. *Am J Obstet Gynecol* 1996 Jun;174(6):1885–93;discussion 1893–5.
- 111 Metules TJ. Are you at risk for spontaneous abortion? *RN: Nat Mag Nursing*. 2000 Aug; 63(8):69–72.
- 112 Meyer BA, Daling JR. Activity level of mother's usual occupation and low infant birth weight. *J Occup Med* 1985 Nov;27(11):841–7.
- 113 Moutquin JM. Socio-economic and psychosocial factors in the management and prevention of preterm labour. *BJOG* 2003 Apr;110 Suppl 20:56–60.
- 114 Mozurkewich EL, Luke B, Avni M, Wolf FM. Working conditions and adverse pregnancy outcome: a meta-analysis. *Obstet Gynecol* 2000 Apr;95(4):623–35.
- 115 Naeye RL, Peters EC. Working during pregnancy: effects on the foetus. *Paediatrics* 1982 Jun; 69(6): 724–7.
- 116 Newman RB, Goldenberg RL, Moawad AH, Iams JD, *et al.* Occupational fatigue and preterm premature rupture of membranes. National Institute of Child Health and Human Development Maternal-Fetal Medicine, Units Network. *Am J Obstet Gynecol* 2001 Feb;184(3):438–46.
- 117 Nicholson PJ, D'Auria DA. Shift work, health, the working time regulations and health assessments. *Occup Med (Lond)*. 1999 Apr;49(3):127–37.
- 118 Nurminen T. Nonstandard working hours and reproductive outcome. *Tyoe Ja Ihminen* 1998; 12(3): 167–176.
- 119 Nurminen T. Shift work and reproductive health. *Scand J Work Environ Health* 1998;24 Suppl 3:28–34.
- 120 Nurminen T. Shift work, foetal development and course of pregnancy. *Scand J Work Environ Health* 1989 Dec;15(6):395–403.
- 121 Nurminen T, Kurppa K, Holmberg PC. Effects of shift work on teratogenic risk and course of pregnancy. *Scandinavian J Work Environ Health* 1987 Apr;13(2):159–160.
- 122 Nurminen T, Lusa S, Ilmarinen J, Kurppa K. Physical work load, foetal development and course of pregnancy. *Scand J Work Environ Health* 1989 Dec;15(6):404–14.
- 123 Osborn LM, Harris DL, Reading JC, Prather MB. Outcome of pregnancies experienced during residency. *J Fam Pract* 1990 Dec;31(6):618–22.
- 124 Oths KS, Dunn LL, Palmer NS. A prospective study of psychosocial job strain and birth outcomes. *Epidemiology* 2001 Nov;12(6):744–6.
- 125 Peoples-Sheps MD, Siegel E, Suchindran CM, Origasa H *et al.* Characteristics of maternal employment during pregnancy: effects on low birthweight. *Am J Public Health* 1991 Aug;81(8):1007–12.
- 126 Perkins LA. Is the night shift worth the risk? *RN: Nat Mag Nursing* 2001 Aug; 64(8):65.
- 127 Baker PN, Luesley D, Drife J. *Obstetrics and Gynaecology: An Evidence-based Text for MRCOG. Gynaecology & Obstetrics.* ISSN: 0340808756.
- 128 Pinhas-Hamiel O, Rotstein Z, Achiron A, Gabbay U *et al.* Pregnancy during residency – an Israeli survey of women physicians. *Health Care Women Int* 1999 Jan–1999 Feb 28;20(1):63–70.
- 129 Pompeii LA. Occupational physical exertion during pregnancy and the risk of preterm delivery and small-for-gestational-age birth. *Dissertations Abstracts International*. 2002;6308B3679–3860.
- 130 Rabkin CS, Anderson HR, Bland JM, Brooke OG *et al.* Maternal activity and birth weight: a prospective, population-based study. *Am J Epidemiol* 1990 Mar;131(3):522–31.
- 131 Ramirez G, Grimes RM, Annegers J F, Davis BR, Slater C H. Occupational physical activity and other risk factors for preterm birth among US Army primigravidas. *Am J Public Health* 1990 Jun; 80(6): 728–30.

- 132 Reilly K. Nutrition, exercise, work, and sex in pregnancy. *Prim Care* 2000 Mar;27(1):105–15.
- 133 Ritsmitchai S, Geater AF, Chonsuviwatvong V. Prolonged standing and physical exertion at work during pregnancy increases the risk of preterm birth for Thai mothers. *J Occupational Health* vol 39 (no 3):217–222.
- 134 Royal College of Physicians. National Sentinel Stroke Audit Report 2004. Prepared on behalf of the Intercollegiate Stroke Working Party. London. RCP, 2005
- 135 Russell R, Reynolds F. Back pain, pregnancy, and childbirth. *BMJ* 1997 Apr 12;314(7087):1062–3.
- 136 Rylander L, Kallen B. Reproductive outcomes among hairdressers. *Scand J Work Environ Health* 2005 Jun;31(3):212–7.
- 137 Saftlas AF, Logsdon-Sackett N, Wang W, Woolson R, Bracken MB. Work, leisure-time physical activity, and risk of pre-eclampsia and gestational hypertension. *Am J Epidemiol* 2004 Oct 15; 160(8):758–65.
- 138 Saurel-Cubizolles MJ, Kaminski M. Pregnant women's working conditions and their changes during pregnancy: a national study in France. *Br J Ind Med* 1987 Apr;44(4):236–43.
- 139 Saurel-Cubizolles MJ, Kaminski M, Llado-Arkipoff J, Du Mazaubrun C *et al.* Pregnancy and its outcome among hospital personnel according to occupation and working conditions. *J Epidemiol Community Health* 1985 Jun;39(2):129–34.
- 140 Saurel-Cubizolles MJ, Zeitlin J, Lelong N, Papiernik E *et al.* Employment, working conditions, and preterm birth: results from the Europop case-control survey. *J Epidemiol Community Health* 2004 May; 58(5):395–401.
- 141 Saurel-Cubizolles MJ, Subtil D, Kaminski M. Is preterm delivery still related to physical working conditions in pregnancy? *J Epidemiol Community Health* (Mar; 45(1):29–34).
- 142 Savitz DA, Brett KM, Dole N, Tse CK. Male and female occupation in relation to miscarriage and preterm delivery in central North Carolina. *Ann Epidemiol.* 1997 Oct;7(7):509–16.
- 143 Savitz DA, Olshan AF, Gallagher K. Maternal occupation and pregnancy outcome. *Epidemiology* 1996 May;7(3):269–74.
- 144 Schneider KT, Deckardt R. The implication of upright posture on pregnancy. *J Perinat Med* 1991; 19(1–2):121–31.
- 145 Scott AJ, LaDou, J. Shiftwork: effects on sleep and health with recommendations for medical surveillance and screening. *Occup Med* 1990 Apr–1990 Jun 30;5(2):273–99.
- 146 Seguin RE. Pregnancy and the working woman: a review. *J Ark Med Soc* 1998 Aug;95(3):115–8.
- 147 Senturia KD. A woman's work is never done: women's work and pregnancy outcome in Albania. *Med Anthropol Q* 1997 Sep;11(3):375–95.
- 148 Sever LE. Ethics, society and occupational reproductive hazards: foetal consequences? *Women Health* 2000;30(3):25–37.
- 149 Shaw GM. Strenuous work, nutrition and adverse pregnancy outcomes: a brief review. *J Nutr* 2003 May;133(5 Suppl 2):1718S–1721S.
- 150 Simpson JL. Are physical activity and employment related to preterm birth and low birth weight? *Am J Obstet Gynecol* 1993 Apr;168(4):1231–8.
- 151 Smith J. Pregnant doctors: health and safety risks in the real world. *BMJ Careers* 2004 Apr.
- 152 Spinillo A, Capuzzo E, Baltaro F, Piazza G, *et al.* The effect of work activity in pregnancy on the risk of foetal growth retardation. *Acta Obstet Gynecol Scand* 1996 Jul;75(6):531–6; ISSN: 0001–6349
- 153 Spinillo A, Capuzzo E, Colonna L, Piazzini G *et al.* The effect of work activity in pregnancy on the risk of severe pre-eclampsia. *Aust N Z J Obstet Gynaecol* 1995 Nov;35(4):380–5.
- 154 Stinson JC, Lee KA. Premature labor and birth: influence of rank and perception of fatigue in active duty military women. *Mil Med* 2003 May;168(5):385–90.
- 155 Strand K, Wergeland E, Bjerkedal T. Work load, job control and risk of leaving work by sickness certification before delivery, Norway 1989. *Scand J Soc Med* 1997 Sep;25(3):193–201.
- 156 Tapp LM. Pregnancy & ERGONOMICS: POTENTIAL HAZARDS & KEY SAFEGUARDS. *Professional Safety Journal* 2000 Aug.
- 157 Taskinen H, Anttila A, Lindbohm ML, Sallmen M, Hemminki K. Spontaneous abortions and congenital malformations among the wives of men occupationally exposed to organic solvents. *Scand J Work Environ Health* 1989 Oct;15(5):345–52.

- 158 Taskinen H, Kyyronen P, Hemminki K. Effects of ultrasound, shortwaves, and physical exertion on pregnancy outcome in physiotherapists. *J Epidemiol Community Health* 1990 Sep; 44(3):196-201.
- 159 Teitelman AM, Welch LS, Hellenbrand KG, Bracken MB. Effect of maternal work activity on preterm birth and low birth weight. *Am J Epidemiol* 1990 Jan;(1):104-13.
- 160 Tophoj A. Assessment of occupational environment of pregnant women by general practitioners and specialists in occupational medicine. *Ugeskr Laeger* 1996 Sep 6;161(36):5019-23.
- 161 Tuntiseranee P, Geater A, Chongsuvivatwong V, Kor-anantakul O. The effect of heavy maternal workload on fetal growth retardation and preterm delivery. A study among southern Thai women. *J Occup Environ Med* 1998 Nov;40(11):1013-21.
- 162 Vegna F, Allegri F, Barba A, Di Giovanni M *et al*. Italian. Relationship between miscarriage and occupation: A survey. *Giornale Italiano Di Medicina Del Lavoro Ed Ergonomia* 2004;Volume XXVI(4/suppl.).
- 163 Victora CG, Adair L, Fall C, Hallal PC, Martorell R, Richter L, Sachdev HPS and the Maternal and Child Undernutrition Study Group. Maternal and child undernutrition: consequences for adult health and human capital. *Lancet* 2008;371:340-357.
- 164 Walker SP, Higgins JR, Permezel M, Brennecke SP, Phil, D. Maternal work and pregnancy. *Aust N Z J Obstet Gynaecol* 1999 May;39(2):144-51.
- 165 Wergeland E, Strand K. Working conditions and prevalence of pre-eclampsia, Norway 1989. *Int J Gynaecol Obstet* 1997 Aug;58(2):189-96.
- 166 Wergeland E, Strand K, Bordaahl PE. Strenuous working conditions and birthweight, Norway 1989. *Acta Obstet Gynecol Scand* 1998 Mar;77(3):263-71;ISSN:0001-6349
- 167 Williams C. Pregnancy at work: Health and Safety for the working woman (book review). *Work Employment and Society* 1999 Mar;13(1):151-154.
- 168 Williams. Maternal work increases risk of pre-eclampsia. *Journal of the National Medical Association* vol 94(no 6):420, June 2002.
- 169 Xu X, Ding M, Li B, Christiani DC. Association of rotating shiftwork with preterm births and low birth weight among never smoking women textile workers in China. *Occup Environ Med* 1994 Jul; 51(7): 470-4.
- 170 Zhang H, Bracken MB. Tree-based two-stage risk factor analysis for spontaneous abortion. Tree-based, two-stage risk factor analysis for spontaneous abortion. *Am J Epidemiol* 1996 Nov; 144(10):989-96.
- 171 Zhu JL, Hjollund NH, Andersen AM, Olsen, J. Shift work, job stress, and late fetal loss: The National Birth Cohort in Denmark. *J Occup Environ Med* 2004 Nov;46(11):1144-9.
- 172 Zhu JL, Hjollund N H, Olsen J. Shift work, duration of pregnancy, and birth weight: the National Birth Cohort in Denmark. *Am J Obstet Gynecol* 2004 Jul;191(1):285-91.
- 173 Zuckerman BS, Frank DA, Hingson R, Morelock S, Kayne HL. Impact of maternal work outside the home during pregnancy on neonatal outcome. *Pediatrics* 1986 Apr;77(4):459-64.

Further copies of these guidelines are available from NHS Plus:
Email: nhsplus@nhs.net

ISBN 978-1-86016-353-1

NHS Plus
Email: nhsplus@nhs.net

www.nhsplus.nhs.uk



Occupational Health Clinical Effectiveness Unit
Royal College of Physicians of London
11 St Andrews Place, Regent's Park, London NW1 4LE

www.rcplondon.ac.uk

OCCUPATIONAL
HEALTH CLINICAL
EFFECTIVENESS UNIT



**Royal College
of Physicians**
Setting higher medical standards

Supported by:

**The Faculty of Occupational Medicine
of the Royal College of Physicians**

6 St Andrews Place, Regent's Park, London NW1 4LB

www.facocmed.ac.uk

fom
Faculty of Occupational Medicine

